

Cdx2 Polyclonal Antibody

Catalog # AP69029

Specification

Cdx2 Polyclonal Antibody - Product Information

Application WB
Primary Accession O99626

Reactivity Human, Mouse, Rat

Host Rabbit Clonality Polyclonal

Cdx2 Polyclonal Antibody - Additional Information

Gene ID 1045

Other Names

CDX2; CDX3; Homeobox protein CDX-2; CDX-3; Caudal-type homeobox protein 2

Dilution

WB~~Western Blot: 1/500 - 1/2000. ELISA: 1/5000. Not yet tested in other applications.

Format

Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.09% (W/V) sodium azide.

Storage Conditions

-20°C

Cdx2 Polyclonal Antibody - Protein Information

Name CDX2

Synonyms CDX3

Function

Transcription factor which regulates the transcription of multiple genes expressed in the intestinal epithelium (By similarity). Binds to the promoter of the intestinal sucrase-isomaltase SI and activates SI transcription (By similarity). Binds to the DNA sequence 5'-ATAAAACTTAT-3' in the promoter region of VDR and activates VDR transcription (By similarity). Binds to and activates transcription of LPH (By similarity). Activates transcription of CLDN2 and intestinal mucin MUC2 (By similarity). Binds to the 5'-AATTTTTTACAACACCT-3' DNA sequence in the promoter region of CA1 and activates CA1 transcription (By similarity). Important in broad range of functions from early differentiation to maintenance of the intestinal epithelial lining of both the small and large intestine. Binds preferentially to methylated DNA (PubMed:28473536).

Cellular Location

Nucleus {ECO:0000250|UniProtKB:P43241}.



Tel: 858.875.1900 Fax: 858.875.1999

Tissue Location


Detected in small intestine, colon and pancreas.

Cdx2 Polyclonal Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- Western Blot
- Blocking Peptides
- Dot Blot
- <u>Immunohistochemistry</u>
- Immunofluorescence
- <u>Immunoprecipitation</u>
- Flow Cytomety
- Cell Culture

Cdx2 Polyclonal Antibody - Images



Cdx2 Polyclonal Antibody - Background

Involved in the transcriptional regulation of multiple genes expressed in the intestinal epithelium. Important in broad range of functions from early differentiation to maintenance of the intestinal epithelial lining of both the small and large intestine. Binds preferentially to methylated DNA (PubMed:28473536).