

Claudin-1 Polyclonal Antibody

Catalog # AP69125

Specification

Claudin-1 Polyclonal Antibody - Product Information

Application
Primary Accession

Reactivity Human, Mouse, Rat Rabbit

Host Rabbit Clonality Polyclonal

Claudin-1 Polyclonal Antibody - Additional Information

Gene ID 9076

Other Names

CLDN1; CLD1; SEMP1; Claudin-1; Senescence-associated epithelial membrane protein

Dilution

WB~~Western Blot: 1/500 - 1/2000. Immunohistochemistry: 1/100 - 1/300. Immunofluorescence:

WB, IHC-P, IF

095832

1/200 - 1/1000. ELISA: 1/40000. Not yet tested in other applications.

IHC-P~~N/A IF~~1:50~200

Format

Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.09% (W/V) sodium azide.

Storage Conditions

-20°C

Claudin-1 Polyclonal Antibody - Protein Information

Name CLDN1

Synonyms CLD1, SEMP1

Function

Claudins function as major constituents of the tight junction complexes that regulate the permeability of epithelia. While some claudin family members play essential roles in the formation of impermeable barriers, others mediate the permeability to ions and small molecules. Often, several claudin family members are coexpressed and interact with each other, and this determines the overall permeability. CLDN1 is required to prevent the paracellular diffusion of small molecules through tight junctions in the epidermis and is required for the normal barrier function of the skin. Required for normal water homeostasis and to prevent excessive water loss through the skin, probably via an indirect effect on the expression levels of other proteins, since CLDN1 itself seems to be dispensable for water barrier formation in keratinocyte tight junctions (PubMed:>23407391/a>).



Cellular Location

Cell junction, tight junction. Cell membrane; Multi-pass membrane protein. Basolateral cell membrane Note=Associates with CD81 and the CLDN1-CD81 complex localizes to the basolateral cell membrane.

Tissue Location


Strongly expressed in liver and kidney. Expressed in heart, brain, spleen, lung and testis.

Claudin-1 Polyclonal Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.


- Western Blot
- Blocking Peptides
- Dot Blot
- Immunohistochemistry
- Immunofluorescence
- <u>Immunoprecipitation</u>
- Flow Cytomety
- Cell Culture

Claudin-1 Polyclonal Antibody - Images



Western Blot analysis of various cells using Claudin-1 Polyclonal Antibody diluted at 1□1000





Claudin-1 Polyclonal Antibody - Background

Claudins function as major constituents of the tight junction complexes that regulate the permeability of epithelia. While some claudin family members play essential roles in the formation of impermeable barriers, others mediate the permeability to ions and small molecules. Often, several claudin family members are coexpressed and interact with each other, and this determines the overall permeability. CLDN1 is required to prevent the paracellular diffusion of small molecules through tight junctions in the epidermis and is required for the normal barrier function of the skin. Required for normal water homeostasis and to prevent excessive water loss through the skin, probably via an indirect effect on the expression levels of other proteins, since CLDN1 itself seems to be dispensable for water barrier formation in keratinocyte tight junctions (PubMed:23407391).