

Claudin-19 Polyclonal Antibody

Catalog # AP69129

Specification

Claudin-19 Polyclonal Antibody - Product Information

Application WB
Primary Accession Q8N6F1
Reactivity Human, Rat
Host Rabbit
Clonality Polyclonal

Claudin-19 Polyclonal Antibody - Additional Information

Gene ID 149461

Other Names CLDN19; Claudin-19

Dilution

WB~~Western Blot: 1/500 - 1/2000. ELISA: 1/10000. Not yet tested in other applications.

Format

Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.09% (W/V) sodium azide.

Storage Conditions -20°C

Claudin-19 Polyclonal Antibody - Protein Information

Name CLDN19 {ECO:0000303|PubMed:25555744, ECO:0000312|HGNC:HGNC:2040}

Function

Forms paracellular channels: coassembles with CLDN16 into tight junction strands with cation-selective channels through the strands, conveying epithelial permeability in a process known as paracellular tight junction permeability (PubMed:18188451, PubMed:28028216). Involved in the maintenance of ion gradients along the nephron. In the thick ascending limb (TAL) of Henle's loop, facilitates sodium paracellular permeability from the interstitial compartment to the lumen, contributing to the lumen-positive transepithelial potential that drives paracellular magnesium and calcium reabsorption (By similarity) (PubMed:17033971, PubMed:25555744). Forms paracellular barriers on its own. In the peripheral nervous system, represents a major constituent of the tight junctions in Schwann cells and contributes to electrical sealing. During retinal neurogenesis, may regulate the barrier properties of tight junctions in retinal pigment epithelium, required for proper retinal tissue differentiation and vision (By similarity) (PubMed:17033971, PubMed:<a href="http://www.uniprot.org/citations/30937396"



target="_blank">30937396).

Cellular Location


Cell junction, tight junction. Cell membrane; Multi-pass membrane protein. Note=Cotrafficks with CLDN16 from ER to tight junctions. Colocalizes with CLDN16 and CLDN3 in cell- cell contact areas of the TAL spatially separated from CLDN10b paracellular channels.

Claudin-19 Polyclonal Antibody - Protocols


Provided below are standard protocols that you may find useful for product applications.

- Western Blot
- Blocking Peptides
- Dot Blot
- <u>Immunohistochemistry</u>
- Immunofluorescence
- Immunoprecipitation
- Flow Cytomety
- Cell Culture

Claudin-19 Polyclonal Antibody - Images







Claudin-19 Polyclonal Antibody - Background

Plays a major role in tight junction-specific obliteration of the intercellular space, through calcium-independent cell-adhesion activity.