

CNOT7 Polyclonal Antibody
Catalog # AP69185**Specification****CNOT7 Polyclonal Antibody - Product Information**

Application	WB
Primary Accession	Q9UIV1
Reactivity	Human, Mouse
Host	Rabbit
Clonality	Polyclonal

CNOT7 Polyclonal Antibody - Additional Information**Gene ID** 29883**Other Names**

CNOT7; CAF1; CCR4-NOT transcription complex subunit 7; BTG1-binding factor 1; CCR4-associated factor 1; CAF-1

Dilution

WB~~Western Blot: 1/500 - 1/2000. Immunohistochemistry: 1/100 - 1/300. ELISA: 1/20000. Not yet tested in other applications.

Format

Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.09% (W/V) sodium azide.

Storage Conditions

-20°C

CNOT7 Polyclonal Antibody - Protein Information**Name** CNOT7**Synonyms** CAF1**Function**

Has 3'-5' poly(A) exoribonuclease activity for synthetic poly(A) RNA substrate (PubMed:20634287, PubMed:31439799, PubMed:19276069). Its function seems to be partially redundant with that of CNOT8 (PubMed:19605561). Catalytic component of the CCR4-NOT complex which is one of the major cellular mRNA deadenylases and is linked to various cellular processes including bulk mRNA degradation, miRNA-mediated repression, translational repression during translational initiation and general transcription regulation (PubMed:20634287, PubMed:31439799, PubMed:19276069).

target="_blank">19276069). During miRNA- mediated repression the complex seems also to act as translational repressor during translational initiation (PubMed:20065043). Additional complex functions may be a consequence of its influence on mRNA expression (PubMed:23236473, PubMed:19276069). Associates with members of the BTG family such as TOB1 and BTG2 and is required for their anti- proliferative activity (PubMed:23236473, PubMed:19276069).

Cellular Location


Nucleus. Cytoplasm, P-body {ECO:0000250|UniProtKB:Q60809}. Cytoplasm, Cytoplasmic ribonucleoprotein granule. Note=NANOS2 promotes its localization to P-body (By similarity). Recruited to cytoplasmic ribonucleoprotein membraneless compartments by CAPRIN1, promoting deadenylation of mRNAs (PubMed:31439799) {ECO:0000250|UniProtKB:Q60809, ECO:0000269|PubMed:31439799}

CNOT7 Polyclonal Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

CNOT7 Polyclonal Antibody - Images



CNOT7 Polyclonal Antibody - Background

Has 3'-5' poly(A) exoribonuclease activity for synthetic poly(A) RNA substrate. Its function seems

to be partially redundant with that of CNOT8. Catalytic component of the CCR4-NOT complex which is one of the major cellular mRNA deadenylases and is linked to various cellular processes including bulk mRNA degradation, miRNA-mediated repression, translational repression during translational initiation and general transcription regulation. During miRNA-mediated repression the complex seems also to act as translational repressor during translational initiation. Additional complex functions may be a consequence of its influence on mRNA expression. Associates with members of the BTG family such as TOB1 and BTG2 and is required for their anti- proliferative activity.