

CNT2 Polyclonal Antibody
Catalog # AP69187**Specification**

CNT2 Polyclonal Antibody - Product Information

Application	WB
Primary Accession	O43868
Reactivity	Human, Mouse, Rat
Host	Rabbit
Clonality	Polyclonal

CNT2 Polyclonal Antibody - Additional Information**Gene ID** 9153**Other Names**

SLC28A2; CNT2; Sodium/nucleoside cotransporter 2; Concentrative nucleoside transporter 2; CNT 2; hCNT2; Na(+)/nucleoside cotransporter 2; Sodium-coupled nucleoside transporter 2; Sodium/purine nucleoside co-transporter; SPNT; Solute carrier

Dilution

WB~~Western Blot: 1/500 - 1/2000. Immunohistochemistry: 1/100 - 1/300. ELISA: 1/10000. Not yet tested in other applications.

Format

Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.09% (W/V) sodium azide.

Storage Conditions

-20°C

CNT2 Polyclonal Antibody - Protein Information**Name** SLC28A2**Synonyms** CNT2**Function**

Sodium-dependent and purine-selective transporter (PubMed:10087507, PubMed:9435697). Exhibits the transport characteristics of the nucleoside transport system cif or N1 subtype (N1/cif) (selective for purine nucleosides and uridine) (PubMed:10087507, PubMed:21795683, PubMed:9435697). Plays a critical role in specific uptake and salvage of purine nucleosides in kidney and other tissues (PubMed:9435697). May contribute to regulate the transport of organic compounds in testes across the blood-testis- barrier

(Probable).

Cellular Location

Membrane; Multi-pass membrane protein. Apicolateral cell membrane; Multi-pass membrane protein. Note=Localized to the apicolateral membranes of Sertoli cells and vascular endothelial cells in testis

Tissue Location


Expressed in heart and skeletal muscle followed by liver, kidney, intestine, pancreas, placenta and brain (PubMed:9435697). Weak expression in lung (PubMed:9435697). In testis, primarily localized to the apicolateral membranes of Sertoli cells and vascular endothelial cells, and weakly expressed in Leydig cells, peritubular myoid cells and germ cells (PubMed:35307651)




CNT2 Polyclonal Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

CNT2 Polyclonal Antibody - Images





CNT2 Polyclonal Antibody - Background

Sodium-dependent and purine-selective transporter. Exhibits the transport characteristics of the nucleoside transport system cif or N1 subtype (N1/cif) (selective for purine nucleosides and uridine). Plays a critical role in specific uptake and salvage of purine nucleosides in kidney and other tissues.