

CstF-50 Polyclonal Antibody
Catalog # AP69327**Specification**

CstF-50 Polyclonal Antibody - Product Information

Application	WB
Primary Accession	Q05048
Reactivity	Human, Mouse, Rat
Host	Rabbit
Clonality	Polyclonal

CstF-50 Polyclonal Antibody - Additional Information**Gene ID** 1477**Other Names**

CSTF1; Cleavage stimulation factor subunit 1; CF-1 50 kDa subunit; Cleavage stimulation factor 50 kDa subunit; CSTF 50 kDa subunit; CstF-50

Dilution

WB~~Western Blot: 1/500 - 1/2000. Immunohistochemistry: 1/100 - 1/300. ELISA: 1/20000. Not yet tested in other applications.

Format

Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.09% (W/V) sodium azide.

Storage Conditions

-20°C

CstF-50 Polyclonal Antibody - Protein Information**Name** CSTF1**Function**

One of the multiple factors required for polyadenylation and 3'-end cleavage of mammalian pre-mRNAs (PubMed:10669729). May be responsible for the interaction of CSTF with other factors to form a stable complex on the pre-mRNA (PubMed:10669729).

Cellular Location



Nucleus.

CstF-50 Polyclonal Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

CstF-50 Polyclonal Antibody - Images



CstF-50 Polyclonal Antibody - Background

One of the multiple factors required for polyadenylation and 3'-end cleavage of mammalian pre-mRNAs. May be responsible for the interaction of CSTF with other factors to form a stable complex on the pre-mRNA.