

CYP4B1 Polyclonal Antibody

Catalog # AP69416

Specification

CYP4B1 Polyclonal Antibody - Product Information

Application WB, IHC-P
Primary Accession P13584
Reactivity Human
Host Rabbit
Clonality Polyclonal

CYP4B1 Polyclonal Antibody - Additional Information

Gene ID 1580

Other Names

CYP4B1; Cytochrome P450 4B1; CYPIVB1; Cytochrome P450-HP

Dilution

WB $\sim\sim$ Western Blot: 1/500 - 1/2000. Immunohistochemistry: 1/100 - 1/300. ELISA: 1/40000. Not yet tested in other applications.

IHC-P~~N/A

Format

Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.09% (W/V) sodium azide.

Storage Conditions

-20°C

CYP4B1 Polyclonal Antibody - Protein Information

Name CYP4B1

Function

Cytochromes P450 are a group of heme-thiolate monooxygenases. In liver microsomes, this enzyme is involved in an NADPH-dependent electron transport pathway. It oxidizes a variety of structurally unrelated compounds, including steroids, fatty acids, and xenobiotics.

Cellular Location

Endoplasmic reticulum membrane; Peripheral membrane protein. Microsome membrane; Peripheral membrane protein

Tissue Location

Detected in the liver and lung (at protein level).


CYP4B1 Polyclonal Antibody - Protocols




Provided below are standard protocols that you may find useful for product applications.

- Western Blot
- Blocking Peptides
- Dot Blot
- <u>Immunohistochemistry</u>
- Immunofluorescence
- Immunoprecipitation
- Flow Cytomety
- Cell Culture

CYP4B1 Polyclonal Antibody - Images





CYP4B1 Polyclonal Antibody - Background

Cytochromes P450 are a group of heme-thiolate monooxygenases. In liver microsomes, this enzyme is involved in an NADPH-dependent electron transport pathway. It oxidizes a variety of structurally unrelated compounds, including steroids, fatty acids, and xenobiotics.