

DGK-B Polyclonal Antibody

Catalog # AP69515

### Specification

# DGK-<sup>β</sup> Polyclonal Antibody - Product Information

Application Primary Accession Reactivity Host Clonality WB <u>O9Y6T7</u> Human, Mouse, Rat Rabbit Polyclonal

### DGK-ß Polyclonal Antibody - Additional Information

Gene ID 1607

**Other Names** DGKB; DAGK2; KIAA0718; Diacylglycerol kinase beta; DAG kinase beta; 90 kDa diacylglycerol kinase; Diglyceride kinase beta; DGK-beta

Dilution

WB~~Western Blot: 1/500 - 1/2000. ELISA: 1/40000. Not yet tested in other applications.

**Format** Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.09% (W/V) sodium azide.

**Storage Conditions** -20°C

# DGK-ß Polyclonal Antibody - Protein Information

Name DGKB

Synonyms DAGK2, KIAA0718

#### Function

Diacylglycerol kinase that converts diacylglycerol/DAG into phosphatidic acid/phosphatidate/PA and regulates the respective levels of these two bioactive lipids (PubMed:<a href="http://www.uniprot.org/citations/11719522" target="\_blank">11719522</a>). Thereby, acts as a central switch between the signaling pathways activated by these second messengers with different cellular targets and opposite effects in numerous biological processes (Probable). Has a higher activity with long-chain diacylglycerols like 1,2-di-(9Z-octadecenoyl)-sn-glycerol compared to 1,2-didecanoyl-sn-glycerol (By similarity). Specifically expressed in brain, it regulates neuron-specific morphological changes including neurite branching and neurite spine formation (By similarity).

#### **Cellular Location**

Postsynaptic cell membrane {ECO:0000250|UniProtKB:Q6NS52}; Peripheral membrane protein {ECO:0000250|UniProtKB:Q6NS52}. Cell membrane; Peripheral membrane protein. Cytoplasm



Note=Translocation to the plasma membrane is induced by phorbol esters

### **Tissue Location**


[Isoform 1]: Specifically expressed in brain but also detected in uterus (PubMed:11719522). In adult brain, expressed in the amygdala, caudate nucleus, and hippocampus (PubMed:11719522)

### DGK-β Polyclonal Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- <u>Western Blot</u>
- Blocking Peptides
- Dot Blot
- Immunohistochemistry
- Immunofluorescence
- Immunoprecipitation
- Flow Cytomety
- <u>Cell Culture</u>

# DGK-β Polyclonal Antibody - Images



# DGK-β Polyclonal Antibody - Background

Exhibits high phosphorylation activity for long-chain diacylglycerols.