

DRAK2 Polyclonal Antibody
Catalog # AP69594**Specification**

DRAK2 Polyclonal Antibody - Product Information

Application	WB, IHC-P
Primary Accession	O94768
Reactivity	Human, Mouse, Rat
Host	Rabbit
Clonality	Polyclonal

DRAK2 Polyclonal Antibody - Additional Information**Gene ID** 9262**Other Names**

STK17B; DRAK2; Serine/threonine-protein kinase 17B; DAP kinase-related apoptosis-inducing protein kinase 2

Dilution

WB~~Western Blot: 1/500 - 1/2000. Immunohistochemistry: 1/100 - 1/300. ELISA: 1/40000. Not yet tested in other applications.

IHC-P~~N/A

Format

Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.09% (W/V) sodium azide.

Storage Conditions

-20°C

DRAK2 Polyclonal Antibody - Protein Information**Name** STK17B**Synonyms** DRAK2**Function**

Phosphorylates myosin light chains (By similarity). Acts as a positive regulator of apoptosis.

Cellular Location

Nucleus. Cell membrane. Endoplasmic reticulum-Golgi intermediate compartment.

Note=Colocalizes with STK17B at the plasma membrane.

Tissue Location


Highly expressed in placenta, lung, pancreas. Lower levels in heart, brain, liver, skeletal muscle and kidney

DRAK2 Polyclonal Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

DRAK2 Polyclonal Antibody - Images



DRAK2 Polyclonal Antibody - Background

Phosphorylates myosin light chains (By similarity). Acts as a positive regulator of apoptosis.