

## **E-cadherin Polyclonal Antibody**

**Catalog # AP69636** 

#### **Specification**

## **E-cadherin Polyclonal Antibody - Product Information**

Application WB
Primary Accession P12830
Reactivity Human
Host Rabbit
Clonality Polyclonal

## E-cadherin Polyclonal Antibody - Additional Information

### Gene ID 999

### **Other Names**

CDH1; CDHE; UVO; Cadherin-1; CAM 120/80; Epithelial cadherin; E-cadherin; Uvomorulin; CD antigen CD324

#### **Dilution**

WB~~Western Blot: 1/500 - 1/2000. Immunohistochemistry: 1/100 - 1/300. Immunofluorescence: 1/200 - 1/1000. ELISA: 1/40000. Not yet tested in other applications.

#### **Format**

Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.09% (W/V) sodium azide.

#### **Storage Conditions**

-20°C

#### E-cadherin Polyclonal Antibody - Protein Information

## Name CDH1

Synonyms CDHE, UVO

### **Function**

Cadherins are calcium-dependent cell adhesion proteins (PubMed:<a href="http://www.uniprot.org/citations/11976333" target="\_blank">11976333</a>). They preferentially interact with themselves in a homophilic manner in connecting cells; cadherins may thus contribute to the sorting of heterogeneous cell types. CDH1 is involved in mechanisms regulating cell-cell adhesions, mobility and proliferation of epithelial cells (PubMed:<a href="http://www.uniprot.org/citations/11976333" target="\_blank">11976333</a>). Has a potent invasive suppressor role. It is a ligand for integrin alpha-E/beta-7.

## **Cellular Location**

Cell junction, adherens junction. Cell membrane; Single-pass type I membrane protein. Endosome. Golgi apparatus, trans-Golgi network. Note=Colocalizes with DLGAP5 at sites of cell-cell contact in intestinal epithelial cells. Anchored to actin microfilaments through association with alpha-, beta-





and gamma-catenin. Sequential proteolysis induced by apoptosis or calcium influx, results in translocation from sites of cell-cell contact to the cytoplasm Colocalizes with RAB11A endosomes during its transport from the Golgi apparatus to the plasma membrane


**Tissue Location**Non-neural epithelial tissues.

# **E-cadherin Polyclonal Antibody - Protocols**


Provided below are standard protocols that you may find useful for product applications.


- Western Blot
- Blocking Peptides
- Dot Blot
- <u>Immunohistochemistry</u>
- Immunofluorescence
- Immunoprecipitation
- Flow Cytomety
- Cell Culture


# E-cadherin Polyclonal Antibody - Images
















# E-cadherin Polyclonal Antibody - Background

Cadherins are calcium-dependent cell adhesion proteins (PubMed:11976333). They preferentially interact with themselves in a homophilic manner in connecting cells; cadherins may thus contribute to the sorting of heterogeneous cell types. CDH1 is involved in mechanisms regulating cell-cell adhesions, mobility and proliferation of epithelial cells (PubMed:11976333). Has a potent invasive suppressor role. It is a ligand for integrin alpha-E/beta-7.