

Factor VIII Polyclonal Antibody
Catalog # AP69836**Specification**

Factor VIII Polyclonal Antibody - Product Information

Application	WB
Primary Accession	P00451
Reactivity	Human, Mouse
Host	Rabbit
Clonality	Polyclonal

Factor VIII Polyclonal Antibody - Additional Information**Gene ID** 2157**Other Names**

F8; F8C; Coagulation factor VIII; Antihemophilic factor; AHF; Procoagulant component

Dilution

WB~~Western Blot: 1/500 - 1/2000. Immunohistochemistry: 1/100 - 1/300. ELISA: 1/10000. Not yet tested in other applications.

Format

Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.09% (W/V) sodium azide.

Storage Conditions

-20°C

Factor VIII Polyclonal Antibody - Protein Information**Name** F8**Synonyms** F8C**Function**

Factor VIII, along with calcium and phospholipid, acts as a cofactor for F9/factor IXa when it converts F10/factor X to the activated form, factor Xa.

Cellular Location

Secreted, extracellular space.



Factor VIII Polyclonal Antibody - Protocols



Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)

- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

Factor VIII Polyclonal Antibody - Images





Factor VIII Polyclonal Antibody - Background

Factor VIII, along with calcium and phospholipid, acts as a cofactor for F9/factor IXa when it converts F10/factor X to the activated form, factor Xa.