

JK (TAOK3) Antibody (C-term)
Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP7012b**Specification**

JK (TAOK3) Antibody (C-term) - Product Information

Application	WB, IHC-P,E
Primary Accession	Q9H2K8
Other Accession	Q6DD27 , Q53UA7 , Q8BYC6 , Q9I9E0
Reactivity	Human
Predicted	Chicken, Mouse, Rat, Xenopus
Host	Rabbit
Clonality	Polyclonal
Isotype	Rabbit IgG
Calculated MW	105406
Antigen Region	671-700

JK (TAOK3) Antibody (C-term) - Additional Information**Gene ID** 51347**Other Names**

Serine/threonine-protein kinase TAO3, Cutaneous T-cell lymphoma-associated antigen HD-CL-09, CTCL-associated antigen HD-CL-09, Dendritic cell-derived protein kinase, JNK/SAPK-inhibitory kinase, Jun kinase-inhibitory kinase, Kinase from chicken homolog A, hKFC-A, Thousand and one amino acid protein 3, TAOK3, DPK, JIK, KDS, MAP3K18

Target/Specificity

This JK (TAOK3) antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 671-700 amino acids from the C-terminal region of human JK (TAOK3).

Dilution

WB~~1:1000
IHC-P~~1:10~50

Format

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is prepared by Saturated Ammonium Sulfate (SAS) precipitation followed by dialysis against PBS.

Storage

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

JK (TAOK3) Antibody (C-term) is for research use only and not for use in diagnostic or therapeutic procedures.

JK (TAOK3) Antibody (C-term) - Protein Information

Name TAOK3

Synonyms DPK, JIK, KDS, MAP3K18

Function Serine/threonine-protein kinase that acts as a regulator of the p38/MAPK14 stress-activated MAPK cascade and of the MAPK8/JNK cascade. Acts as an activator of the p38/MAPK14 stress-activated MAPK cascade. In response to DNA damage, involved in the G2/M transition DNA damage checkpoint by activating the p38/MAPK14 stress-activated MAPK cascade, probably by mediating phosphorylation of upstream MAP2K3 and MAP2K6 kinases. Inhibits basal activity of MAPK8/JNK cascade and diminishes its activation in response epidermal growth factor (EGF).

Cellular Location

Cytoplasm. Cell membrane; Peripheral membrane protein. Note=Also localized to the peripheral cell membrane

Tissue Location


Ubiquitously expressed at a low level, and highly expressed in peripheral blood leukocytes (PBLs), thymus, spleen, kidney, skeletal muscle, heart and liver.

JIK (TAOK3) Antibody (C-term) - Protocols

Provided below are standard protocols that you may find useful for product applications.


- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

JIK (TAOK3) Antibody (C-term) - Images



Western blot analysis of TAOK3 (arrow) using rabbit polyclonal TAOK3 Antibody (C-term)(Cat.#AP7012b). 293 cell lysates (2 ug/lane) either nontransfected (Lane 1) or transiently

transfected with the TAOK3 gene (Lane 2) (Origene Technologies).



Formalin-fixed and paraffin-embedded human hepatocarcinoma tissue reacted with TAOK3 antibody (C-term)(Cat.#AP7012b), which was peroxidase-conjugated to the secondary antibody, followed by DAB staining. This data demonstrates the use of this antibody for immunohistochemistry; clinical relevance has not been evaluated.

JIK (TAOK3) Antibody (C-term) - Background

TAOK3 is negatively regulated by epidermal growth factor (EGF) and Inhibits the basal activity of Jun kinase. When TAOK3 is overexpressed, it may activate ERK1/ERK2 and JNK/SAPK.

JIK (TAOK3) Antibody (C-term) - References

- Hartmann,T.B.,Br. J. Dermatol. 150 (2), 252-258 (2004)
- Yustein,J.T.,Oncogene 22 (40), 6129-6141 (2003)
- Yoneda,T.,J. Biol. Chem. 276 (17), 13935-13940 (2001)
- Zhang,W.,Biochem. Biophys. Res. Commun. 274 (3), 872-879 (2000)