

HBP1 Polyclonal Antibody

Catalog # AP70286

Specification

HBP1 Polyclonal Antibody - Product Information

Application Primary Accession Reactivity Host Clonality WB, IHC-P <u>O60381</u> Human, Mouse, Rat, Monkey Rabbit Polyclonal

HBP1 Polyclonal Antibody - Additional Information

Gene ID 26959

Other Names HBP1; HMG box-containing protein 1; HMG box transcription factor 1; High mobility group box transcription factor 1

Dilution WB~~Western Blot: 1/500 - 1/2000. Immunohistochemistry: 1/100 - 1/300. ELISA: 1/10000. Not yet tested in other applications. IHC-P~~N/A

Format Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.09% (W/V) sodium azide.

Storage Conditions -20°C

HBP1 Polyclonal Antibody - Protein Information

Name HBP1

Function

Transcriptional repressor that binds to the promoter region of target genes. Plays a role in the regulation of the cell cycle and of the Wnt pathway. Binds preferentially to the sequence 5'-TTCATTCATTCA-3'. Binding to the histone H1.0 promoter is enhanced by interaction with RB1. Disrupts the interaction between DNA and TCF4.

Cellular Location Nucleus {ECO:0000255|PROSITE-ProRule:PRU00267, ECO:0000269|PubMed:10562551}


HBP1 Polyclonal Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.



- <u>Western Blot</u>
- <u>Blocking Peptides</u>
- Dot Blot
- Immunohistochemistry
- <u>Immunofluorescence</u>
- Immunoprecipitation
- Flow Cytomety
- <u>Cell Culture</u>

HBP1 Polyclonal Antibody - Images



HBP1 Polyclonal Antibody - Background

Transcriptional repressor that binds to the promoter region of target genes. Plays a role in the regulation of the cell cycle and of the Wnt pathway. Binds preferentially to the sequence 5'-TTCATTCATTCA-3'. Binding to the H1F0 promoter is enhanced by interaction with RB1. Disrupts the interaction between DNA and TCF4.