

## **hCAP-H Polyclonal Antibody**

**Catalog # AP70289**

### **Specification**

#### **hCAP-H Polyclonal Antibody - Product Information**

Application	WB
Primary Accession	<a href="#">Q15003</a>
Reactivity	Human
Host	Rabbit
Clonality	Polyclonal

#### **hCAP-H Polyclonal Antibody - Additional Information**

##### **Gene ID 23397**

##### **Other Names**

NCAPH; BRRN; BRRN1; CAPH; KIAA0074; Condensin complex subunit 2; Barren homolog protein 1; Chromosome-associated protein H; hCAP-H; Non-SMC condensin I complex subunit H; XCAP-H homolog

##### **Dilution**

WB~~Western Blot: 1/500 - 1/2000. ELISA: 1/40000. Not yet tested in other applications.

##### **Format**

Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.09% (W/V) sodium azide.

##### **Storage Conditions**

-20°C

#### **hCAP-H Polyclonal Antibody - Protein Information**

**Name** NCAPH {ECO:0000303|PubMed:27737959, ECO:0000312|HGNC:HGNC:1112}

##### **Function**

Regulatory subunit of the condensin complex, a complex required for conversion of interphase chromatin into mitotic-like condensed chromosomes. The condensin complex probably introduces positive supercoils into relaxed DNA in the presence of type I topoisomerases and converts nicked DNA into positive knotted forms in the presence of type II topoisomerases (PubMed:<a href="http://www.uniprot.org/citations/11136719" target="\_blank">11136719</a>). Early in neurogenesis, may play an essential role to ensure accurate mitotic chromosome condensation in neuron stem cells, ultimately affecting neuron pool and cortex size (PubMed:<a href="http://www.uniprot.org/citations/27737959" target="\_blank">27737959</a>).

##### **Cellular Location**

Nucleus. Cytoplasm. Chromosome. Note=In interphase cells, the majority of the condensin complex is found in the cytoplasm, while a minority of the complex is associated with chromatin. A subpopulation of the complex however remains associated with chromosome foci in interphase cells. During mitosis, most of the condensin complex is associated with the chromatin. At the onset

of prophase, the regulatory subunits of the complex are phosphorylated by CDK1, leading to condensin's association with chromosome arms and to chromosome condensation. Dissociation from chromosomes is observed in late telophase


**Tissue Location**

Widely expressed at low level. Expressed in proliferating cells.

**hCAP-H Polyclonal Antibody - Protocols**

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

**hCAP-H Polyclonal Antibody - Images**

Western Blot analysis of COLO cells using hCAP-H Polyclonal Antibody

**hCAP-H Polyclonal Antibody - Background**

Regulatory subunit of the condensin complex, a complex required for conversion of interphase chromatin into mitotic-like condensed chromosomes. The condensin complex probably introduces positive supercoils into relaxed DNA in the presence of type I topoisomerases and converts nicked DNA into positive knotted forms in the presence of type II topoisomerases (PubMed:11136719). Early in neurogenesis, may play an essential role to ensure accurate mitotic chromosome condensation in neuron stem cells, ultimately affecting neuron pool and cortex size (PubMed:27737959).