

# Histamine H3 Receptor Polyclonal Antibody

**Catalog # AP70325** 

## Specification

## **Histamine H3 Receptor Polyclonal Antibody - Product Information**

**Application Primary Accession** 

Reactivity Host Clonality

WB, IHC-P, IF **Q9Y5N1** 

Human, Mouse, Rat

Rabbit **Polyclonal** 

## Histamine H3 Receptor Polyclonal Antibody - Additional Information

Gene ID 11255

#### **Other Names**

HRH3; GPCR97; Histamine H3 receptor; H3R; HH3R; G-protein coupled receptor 97

#### Dilution

WB~~Western Blot: 1/500 - 1/2000. Immunohistochemistry: 1/100 - 1/300. Immunofluorescence: 1/200 - 1/1000. ELISA: 1/5000. Not yet tested in other applications. IHC-P~~N/A

IF~~1:50~200

### **Format**

Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.09% (W/V) sodium azide.

## **Storage Conditions**

-20°C

## **Histamine H3 Receptor Polyclonal Antibody - Protein Information**

### Name HRH3

Synonyms GPCR97

#### **Function**

The H3 subclass of histamine receptors could mediate the histamine signals in CNS and peripheral nervous system. Signals through the inhibition of adenylate cyclase and displays high constitutive activity (spontaneous activity in the absence of agonist). Agonist stimulation of isoform 3 neither modified adenylate cyclase activity nor induced intracellular calcium mobilization.

#### **Cellular Location**

Cell membrane; Multi-pass membrane protein.

### **Tissue Location**

Expressed predominantly in the CNS, with the greatest expression in the thalamus and caudate nucleus. The various isoforms are mainly coexpressed in brain, but their relative expression level




varies in a region-specific manner. Isoform 3 and isoform 7 are highly expressed in the thalamus, caudate nucleus and cerebellum while isoform 5 and isoform 6 show a poor expression. Isoform 5 and isoform 6 show a high expression in the amygdala, substantia nigra, cerebral cortex and hypothalamus. Isoform 7 is not found in hypothalamus or substantia nigra

## **Histamine H3 Receptor Polyclonal Antibody - Protocols**

Provided below are standard protocols that you may find useful for product applications.

- Western Blot
- Blocking Peptides
- Dot Blot
- Immunohistochemistry
- Immunofluorescence
- Immunoprecipitation
- Flow Cytomety
- Cell Culture

## **Histamine H3 Receptor Polyclonal Antibody - Images**



Western Blot analysis of COLO205 cells using Histamine H3 Receptor Polyclonal Antibody

# Histamine H3 Receptor Polyclonal Antibody - Background

The H3 subclass of histamine receptors could mediate the histamine signals in CNS and peripheral nervous system. Signals through the inhibition of adenylate cyclase and displays high constitutive activity (spontaneous activity in the absence of agonist). Agonist stimulation of isoform 3 neither modified adenylate cyclase activity nor induced intracellular calcium mobilization.