

c-AMPProtein Kinase Catalytic subunit Antibody (N-term E32)
Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP7047d

Specification

c-AMPProtein Kinase Catalytic subunit Antibody (N-term E32) - Product Information

Application	WB,E
Primary Accession	P22694
Reactivity	Human
Host	Rabbit
Clonality	Polyclonal
Isotype	Rabbit IgG
Antigen Region	17-47

c-AMPProtein Kinase Catalytic subunit Antibody (N-term E32) - Additional Information

Gene ID 5567

Other Names

cAMP-dependent protein kinase catalytic subunit beta, PKA C-beta, PRKACB

Target/Specificity

This c-AMPProtein Kinase Catalytic subunit antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 17-47 amino acids from the N-terminal region of human c-AMPProtein Kinase Catalytic subunit.

Dilution

WB~~1:1000

Format

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is purified through a protein A column, followed by peptide affinity purification.

Storage

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

c-AMPProtein Kinase Catalytic subunit Antibody (N-term E32) is for research use only and not for use in diagnostic or therapeutic procedures.

c-AMPProtein Kinase Catalytic subunit Antibody (N-term E32) - Protein Information

Name PRKACB

Function Mediates cAMP-dependent signaling triggered by receptor binding to GPCRs (PubMed:[12420224](#), PubMed:[21423175](#), PubMed:[31112131](#)). PKA activation regulates diverse cellular processes such as cell proliferation, the cell cycle, differentiation and regulation of

microtubule dynamics, chromatin condensation and decondensation, nuclear envelope disassembly and reassembly, as well as regulation of intracellular transport mechanisms and ion flux (PubMed:[12420224](#), PubMed:[21423175](#)). Regulates the abundance of compartmentalized pools of its regulatory subunits through phosphorylation of PJA2 which binds and ubiquitinates these subunits, leading to their subsequent proteolysis (PubMed:[12420224](#), PubMed:[21423175](#)). Phosphorylates GPKOW which regulates its ability to bind RNA (PubMed:[21880142](#)). Acts as a negative regulator of mTORC1 by mediating phosphorylation of RPTOR (PubMed:[31112131](#)).

Cellular Location

Cytoplasm. Cell membrane. Membrane; Lipid- anchor. Nucleus {ECO:0000250|UniProtKB:P05131} Note=Translocates into the nucleus (monomeric catalytic subunit). The inactive holoenzyme is found in the cytoplasm {ECO:0000250|UniProtKB:P05131}

Tissue Location


Isoform 1 is most abundant in the brain, with low level expression in kidney. Isoform 2 is predominantly expressed in thymus, spleen and kidney. Isoform 3 and isoform 4 are only expressed in the brain.

c-AMPProtein Kinase Catalytic subunit Antibody (N-term E32) - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

c-AMPProtein Kinase Catalytic subunit Antibody (N-term E32) - Images



The anti-PRKACB Pab (Cat. #AP7047d) is used in Western blot to detect PRKACB in A375 cell lysate.

c-AMPProtein Kinase Catalytic subunit Antibody (N-term E32) - Background

cAMP is a signaling molecule important for a variety of cellular functions. cAMP exerts its effects by activating the cAMP-dependent protein kinase (AMPK), which transduces the signal through phosphorylation of different target proteins. The inactive holoenzyme of AMPK is a tetramer composed of two regulatory and two catalytic subunits. cAMP causes the dissociation of the inactive holoenzyme into a dimer of regulatory subunits bound to four cAMP and two free monomeric catalytic subunits. Four different regulatory subunits and three catalytic subunits of AMPK have been identified in humans. PRKACB is a member of the Ser/Thr protein kinase family and is a catalytic subunit of AMPK.

c-AMPProtein Kinase Catalytic subnunit Antibody (N-term E32) - References

- Dwivedi, Y., et al., Biol. Psychiatry 55(3):234-243 (2004).
Cartier, C., et al., J. Biol. Chem. 278(37):35211-35219 (2003).
Higuchi, H., et al., EMBO J. 22(8):1790-1800 (2003).
Wu, K.J., et al., Oncogene 21(51):7872-7882 (2002).
Jiang, C.H., et al., Proc. Natl. Acad. Sci. U.S.A. 98(4):1930-1934 (2001).