

IL-2 Polyclonal Antibody Catalog # AP70506

Specification

IL-2 Polyclonal Antibody - Product Information

Application	WB, IHC-P
Primary Accession	P60568
Reactivity	Human, Rat
Host	Rabbit
Clonality	Polyclonal

IL-2 Polyclonal Antibody - Additional Information

Gene ID 3558

Other Names

IL2; Interleukin-2; IL-2; T-cell growth factor; TCGF; Aldesleukin

Dilution

WB~~1:1000

IHC-P~~N/A

Format

Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.09% (W/V) sodium azide.

Storage Conditions

-20°C

IL-2 Polyclonal Antibody - Protein Information

Name IL2

Function

Cytokine produced by activated CD4-positive helper T-cells and to a lesser extend activated CD8-positive T-cells and natural killer (NK) cells that plays pivotal roles in the immune response and tolerance (PubMed:6438535). Binds to a receptor complex composed of either the high-affinity trimeric IL-2R (IL2RA/CD25, IL2RB/CD122 and IL2RG/CD132) or the low-affinity dimeric IL-2R (IL2RB and IL2RG) (PubMed:16293754, PubMed:16477002). Interaction with the receptor leads to oligomerization and conformation changes in the IL-2R subunits resulting in downstream signaling starting with phosphorylation of JAK1 and JAK3 (PubMed:7973659). In turn, JAK1 and JAK3 phosphorylate the receptor to form a docking site leading to the phosphorylation of several substrates including STAT5 (PubMed:8580378). This process leads to activation of several pathways including STAT, phosphoinositide-3- kinase/PI3K and mitogen-activated protein kinase/MAPK pathways (PubMed:16293754).

[25142963](http://www.uniprot.org/citations/25142963)). Functions as a T-cell growth factor and can increase NK-cell cytolytic activity as well (PubMed:[6608729](http://www.uniprot.org/citations/6608729)). Promotes strong proliferation of activated B-cells and subsequently immunoglobulin production (PubMed:[6438535](http://www.uniprot.org/citations/6438535)). Plays a pivotal role in regulating the adaptive immune system by controlling the survival and proliferation of regulatory T-cells, which are required for the maintenance of immune tolerance. Moreover, participates in the differentiation and homeostasis of effector T-cell subsets, including Th1, Th2, Th17 as well as memory CD8-positive T-cells.

Cellular Location


Secreted.

IL-2 Polyclonal Antibody - Protocols


Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)


IL-2 Polyclonal Antibody - Images




Western Blot analysis of various cells using IL-2 Polyclonal Antibody



Immunohistochemical analysis of paraffin-embedded Human kidney. 1, Antibody was diluted at 1:100(4°,overnight). 2, High-pressure and temperature EDTA, pH8.0 was used for antigen retrieval. 3,Secondary antibody was diluted at 1:200(room temperature, 30min).



Immunohistochemical analysis of paraffin-embedded Human kidney. 1, Antibody was diluted at 1:100(4°,overnight). 2, High-pressure and temperature EDTA, pH8.0 was used for antigen retrieval. 3,Secondary antibody was diluted at 1:200(room temperature, 30min).



Immunohistochemical analysis of paraffin-embedded Human kidney. 1, Antibody was diluted at 1:100(4°,overnight). 2, High-pressure and temperature EDTA, pH8.0 was used for antigen retrieval. 3,Secondary antibody was diluted at 1:200(room temperature, 30min).

IL-2 Polyclonal Antibody - Background

Produced by T-cells in response to antigenic or mitogenic stimulation, this protein is required for T-cell proliferation and other activities crucial to regulation of the immune response. Can stimulate B-cells, monocytes, lymphokine- activated killer cells, natural killer cells, and glioma cells.