

IRTKS Polyclonal Antibody
Catalog # AP70594**Specification**

IRTKS Polyclonal Antibody - Product Information

Application	WB, IHC-P
Primary Accession	Q9UHR4
Reactivity	Human, Mouse, Rat
Host	Rabbit
Clonality	Polyclonal

IRTKS Polyclonal Antibody - Additional Information**Gene ID** 55971**Other Names**BAIAP2L1; IRTKS; Brain-specific angiogenesis inhibitor 1-associated protein 2-like protein 1;
BAI1-associated protein 2-like protein 1; Insulin receptor tyrosine kinase substrate**Dilution**

WB~~Western Blot: 1/500 - 1/2000. Immunohistochemistry: 1/100 - 1/300. ELISA: 1/5000. Not yet tested in other applications.

IHC-P~~N/A

Format

Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.09% (W/V) sodium azide.

Storage Conditions

-20°C

IRTKS Polyclonal Antibody - Protein Information**Name** BAIAP2L1**Synonyms** IRTKS**Function**

May function as adapter protein. Involved in the formation of clusters of actin bundles. Plays a role in the reorganization of the actin cytoskeleton in response to bacterial infection.

Cellular Location



Cytoplasm, cytoskeleton. Note=Recruited to actin pedestals that are formed upon infection by bacteria at bacterial attachment sites

IRTKS Polyclonal Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

IRTKS Polyclonal Antibody - Images



IRTKS Polyclonal Antibody - Background

May function as adapter protein. Involved in the formation of clusters of actin bundles. Plays a role in the reorganization of the actin cytoskeleton in response to bacterial infection.