

Laminin γ -3 Polyclonal Antibody
Catalog # AP70715**Specification**

Laminin γ -3 Polyclonal Antibody - Product Information

Application	WB, IHC-P, IF
Primary Accession	Q9Y6N6
Reactivity	Human, Mouse
Host	Rabbit
Clonality	Polyclonal

Laminin γ -3 Polyclonal Antibody - Additional Information**Gene ID** 10319**Other Names**LAMC3; Laminin subunit gamma-3; Laminin-12 subunit gamma; Laminin-14 subunit gamma;
Laminin-15 subunit gamma**Dilution**WB~~Western Blot: 1/500 - 1/2000. Immunohistochemistry: 1/100 - 1/300. Immunofluorescence:
1/200 - 1/1000. ELISA: 1/40000. Not yet tested in other applications.
IHC-P~~N/A
IF~~1:50~200**Format**

Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.09% (W/V) sodium azide.

Storage Conditions

-20°C

Laminin γ -3 Polyclonal Antibody - Protein Information**Name** LAMC3**Function**

Binding to cells via a high affinity receptor, laminin is thought to mediate the attachment, migration and organization of cells into tissues during embryonic development by interacting with other extracellular matrix components.

Cellular Location

Secreted, extracellular space, extracellular matrix, basement membrane

Tissue Location



Broadly expressed in: skin, heart, lung, and the reproductive tracts

Laminin γ -3 Polyclonal Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

Laminin γ -3 Polyclonal Antibody - Images



Laminin γ -3 Polyclonal Antibody - Background

Binding to cells via a high affinity receptor, laminin is thought to mediate the attachment,

migration and organization of cells into tissues during embryonic development by interacting with other extracellular matrix components.