

CKM Antibody (N-term)
Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP7073a**Specification**

CKM Antibody (N-term) - Product Information

Application	WB,E
Primary Accession	P06732
Reactivity	Human, Mouse
Host	Rabbit
Clonality	Polyclonal
Isotype	Rabbit IgG
Calculated MW	43101
Antigen Region	1-30

CKM Antibody (N-term) - Additional Information**Gene ID** 1158**Other Names**

Creatine kinase M-type, Creatine kinase M chain, M-CK, Creatine kinase M-type, N-terminally processed, CKM, CKMM

Target/Specificity

This CKM antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 1-30 amino acids from the N-terminal region of human CKM.

Dilution

WB~~1:1000

E~~Use at an assay dependent concentration.

Format

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is prepared by Saturated Ammonium Sulfate (SAS) precipitation followed by dialysis against PBS.

Storage

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

CKM Antibody (N-term) is for research use only and not for use in diagnostic or therapeutic procedures.

CKM Antibody (N-term) - Protein Information**Name** CKM**Synonyms** CKMM

Function Reversibly catalyzes the transfer of phosphate between ATP and various phosphogens (e.g. creatine phosphate). Creatine kinase isoenzymes play a central role in energy transduction in tissues with large, fluctuating energy demands, such as skeletal muscle, heart, brain and spermatozoa.

Cellular Location


Cytoplasm.

CKM Antibody (N-term) - Protocols


Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

CKM Antibody (N-term) - Images



The anti-CKM Pab (Cat. #AP7073a) is used in Western blot to detect CKM in mouse heart tissue lysate.



Western blot analysis of CKM (arrow) using rabbit polyclonal CKM-T6 (Cat. #AP7073a). 293 cell lysates (2 ug/lane) either nontransfected (Lane 1) or transiently transfected (Lane 2) with the CKM gene.

CKM Antibody (N-term) - Background

Creatine kinase isoenzymes play a central role in energy transduction in tissues with large, fluctuating energy demands, such as skeletal muscle, heart, brain and spermatozoa. The CKM isoform, predominant in skeletal muscle and heart tissue, is a cytoplasmic enzyme involved in energy homeostasis and is an important serum marker for myocardial infarction. CKM reversibly catalyzes the transfer of phosphate between ATP and various phosphogens such as creatine phosphate. It acts as a homodimer in striated muscle as well as in other tissues, and as a heterodimer with a similar brain isozyme in heart. The encoded protein is a member of the ATP:guanido phosphotransferase protein family.