

Med18 Polyclonal Antibody
Catalog # AP70879**Specification**

Med18 Polyclonal Antibody - Product Information

Application	WB
Primary Accession	Q9BUE0
Reactivity	Human, Mouse, Rat
Host	Rabbit
Clonality	Polyclonal

Med18 Polyclonal Antibody - Additional Information**Gene ID** 54797**Other Names**

MED18; Mediator of RNA polymerase II transcription subunit 18; Mediator complex subunit 18; p28b

Dilution

WB~~Western Blot: 1/500 - 1/2000. ELISA: 1/40000. Not yet tested in other applications.

Format

Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.09% (W/V) sodium azide.

Storage Conditions

-20°C

Med18 Polyclonal Antibody - Protein Information**Name** MED18**Function**

Component of the Mediator complex, a coactivator involved in the regulated transcription of nearly all RNA polymerase II-dependent genes. Mediator functions as a bridge to convey information from gene- specific regulatory proteins to the basal RNA polymerase II transcription machinery. Mediator is recruited to promoters by direct interactions with regulatory proteins and serves as a scaffold for the assembly of a functional preinitiation complex with RNA polymerase II and the general transcription factors.

Cellular Location



Nucleus.

Med18 Polyclonal Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

Med18 Polyclonal Antibody - Images



Med18 Polyclonal Antibody - Background

Component of the Mediator complex, a coactivator involved in the regulated transcription of nearly all RNA polymerase II-dependent genes. Mediator functions as a bridge to convey information from gene-specific regulatory proteins to the basal RNA polymerase II transcription machinery. Mediator is recruited to promoters by direct interactions with regulatory proteins and serves as a scaffold for the assembly of a functional preinitiation complex with RNA polymerase II

and the general transcription factors.