

NARG1 Polyclonal Antibody
Catalog # AP71164**Specification****NARG1 Polyclonal Antibody - Product Information**

Application	WB
Primary Accession	Q9BXJ9
Reactivity	Human, Mouse
Host	Rabbit
Clonality	Polyclonal

NARG1 Polyclonal Antibody - Additional Information**Gene ID** 80155**Other Names**

NAA15; GA19; NARG1; NATH; TBDN100; N-alpha-acetyltransferase 15; NatA auxiliary subunit; Gastric cancer antigen Ga19; N-terminal acetyltransferase; NMDA receptor-regulated protein 1; Protein tubedown-1; Tbdn100

Dilution

WB~~Western Blot: 1/500 - 1/2000. Immunohistochemistry: 1/100 - 1/300. Immunofluorescence: 1/200 - 1/1000. ELISA: 1/20000. Not yet tested in other applications.

Format

Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.09% (W/V) sodium azide.

Storage Conditions

-20°C

NARG1 Polyclonal Antibody - Protein Information**Name** NAA15**Synonyms** GA19, NARG1, NATH, TBDN100**Function**

Auxillary subunit of N-terminal acetyltransferase complexes which display alpha (N-terminal) acetyltransferase (NAT) activity (PubMed:15496142, PubMed:20154145, PubMed:29754825, PubMed:32042062). The NAT activity may be important for vascular, hematopoietic and neuronal growth and development (PubMed:15496142). Required to control retinal neovascularization in adult ocular endothelial cells (PubMed:11687548). In complex with XRCC6 and XRCC5 (Ku80), up-regulates transcription from the osteocalcin promoter

(PubMed:12145306).

Cellular Location

Cytoplasm. Nucleus. Note=Mainly cytoplasmic, nuclear in some cases. Present in the free cytosolic and cytoskeleton- bound polysomes, but not in the membrane-bound polysomes

Tissue Location


Expressed at high levels in testis and in ocular endothelial cells. Also found in brain (corpus callosum), heart, colon, bone marrow and at lower levels in most adult tissues, including thyroid, liver, pancreas, mammary and salivary glands, lung, ovary, urogenital system and upper gastrointestinal tract. Overexpressed in gastric cancer, in papillary thyroid carcinomas and in a Burkitt lymphoma cell line (Daudi). Specifically suppressed in abnormal proliferating blood vessels in eyes of patients with proliferative diabetic retinopathy.


NARG1 Polyclonal Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

NARG1 Polyclonal Antibody - Images





NARG1 Polyclonal Antibody - Background

Auxillary subunit of the N-terminal acetyltransferase A (NatA) complex which displays alpha (N-terminal) acetyltransferase activity. The NAT activity may be important for vascular, hematopoietic and neuronal growth and development. Required to control retinal neovascularization in adult ocular endothelial cells. In complex with XRCC6 and XRCC5 (Ku80), up-regulates transcription from the osteocalcin promoter.