

### **NDUFB10 Polyclonal Antibody**

**Catalog # AP71196** 

## **Specification**

### **NDUFB10 Polyclonal Antibody - Product Information**

Application WB, IHC-P
Primary Accession
Reactivity Human
Host Rabbit
Clonality Polyclonal

## **NDUFB10 Polyclonal Antibody - Additional Information**

### **Gene ID 4716**

#### **Other Names**

NDUFB10; NADH dehydrogenase [ubiquinone] 1 beta subcomplex subunit 10; Complex I-PDSW; CI-PDSW; NADH-ubiquinone oxidoreductase PDSW subunit

#### **Dilution**

WB~~Western Blot: 1/500 - 1/2000. Immunohistochemistry: 1/100 - 1/300. ELISA: 1/40000. Not yet tested in other applications. IHC-P~ $\sim$ N/A

### **Format**

Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.09% (W/V) sodium azide.

# **Storage Conditions**

-20°C

## **NDUFB10 Polyclonal Antibody - Protein Information**

### Name NDUFB10

#### **Function**

Accessory subunit that is involved in the functional assembly of the mitochondrial respiratory chain complex I. Complex I has an NADH dehydrogenase activity with ubiquinone as an immediate electron acceptor and mediates the transfer of electrons from NADH to the respiratory chain.

#### **Cellular Location**

Mitochondrion inner membrane; Peripheral membrane protein; Matrix side


## **NDUFB10 Polyclonal Antibody - Protocols**


Provided below are standard protocols that you may find useful for product applications.



- Western Blot
- Blocking Peptides
- Dot Blot
- Immunohistochemistry
- Immunofluorescence
- <u>Immunoprecipitation</u>
- Flow Cytomety
- Cell Culture

# **NDUFB10 Polyclonal Antibody - Images**





# NDUFB10 Polyclonal Antibody - Background

Accessory subunit of the mitochondrial membrane respiratory chain NADH dehydrogenase (Complex I), that is believed not to be involved in catalysis. Complex I functions in the transfer of electrons from NADH to the respiratory chain. The immediate electron acceptor for the enzyme is believed to be ubiquinone.