

NF-YC Polyclonal Antibody

Catalog # AP71275

Specification

NF-YC Polyclonal Antibody - Product Information

Application WB, IHC-P Primary Accession 013952

Reactivity Human, Mouse, Rat

Host Rabbit Clonality Polyclonal

NF-YC Polyclonal Antibody - Additional Information

Gene ID 4802

Other Names

NFYC; Nuclear transcription factor Y subunit gamma; CAAT box DNA-binding protein subunit C; Nuclear transcription factor Y subunit C; NF-YC; Transactivator HSM-1/2

Dilution

WB~~Western Blot: 1/500 - 1/2000. Immunohistochemistry: 1/100 - 1/300. ELISA: 1/10000. Not yet tested in other applications.

IHC-P~~N/A

Format

Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.09% (W/V) sodium azide.

Storage Conditions

-20°C

NF-YC Polyclonal Antibody - Protein Information

Name NFYC

Function


Component of the sequence-specific heterotrimeric transcription factor (NF-Y) which specifically recognizes a 5'-CCAAT-3' box motif found in the promoters of its target genes. NF-Y can function as both an activator and a repressor, depending on its interacting cofactors.

Cellular Location

Nucleus.

NF-YC Polyclonal Antibody - Protocols


Provided below are standard protocols that you may find useful for product applications.







- Western Blot
- Blocking Peptides
- Dot Blot
- <u>Immunohistochemistry</u>
- <u>Immunofluorescence</u>
- <u>Immunoprecipitation</u>
- Flow Cytomety
- Cell Culture

NF-YC Polyclonal Antibody - Images









NF-YC Polyclonal Antibody - Background

Component of the sequence-specific heterotrimeric transcription factor (NF-Y) which specifically recognizes a 5'- CCAAT-3' box motif found in the promoters of its target genes. NF- Y can function as both an activator and a repressor, depending on its interacting cofactors.