

NF-YC Polyclonal Antibody
Catalog # AP71275**Specification**

NF-YC Polyclonal Antibody - Product Information

Application	WB
Primary Accession	Q13952
Reactivity	Human, Mouse, Rat
Host	Rabbit
Clonality	Polyclonal

NF-YC Polyclonal Antibody - Additional Information**Gene ID** 4802**Other Names**

NFYC; Nuclear transcription factor Y subunit gamma; CAAT box DNA-binding protein subunit C; Nuclear transcription factor Y subunit C; NF-YC; Transactivator HSM-1/2

Dilution

WB~~Western Blot: 1/500 - 1/2000. Immunohistochemistry: 1/100 - 1/300. ELISA: 1/10000. Not yet tested in other applications.

Format

Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.09% (W/V) sodium azide.

Storage Conditions

-20°C

NF-YC Polyclonal Antibody - Protein Information**Name** NFYC**Function**

Component of the sequence-specific heterotrimeric transcription factor (NF-Y) which specifically recognizes a 5'-CCAAT-3' box motif found in the promoters of its target genes. NF-Y can function as both an activator and a repressor, depending on its interacting cofactors.

Cellular Location

Nucleus.



NF-YC Polyclonal Antibody - Protocols



Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)

- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

NF-YC Polyclonal Antibody - Images





NF-YC Polyclonal Antibody - Background

Component of the sequence-specific heterotrimeric transcription factor (NF-Y) which specifically recognizes a 5'- CCAAT-3' box motif found in the promoters of its target genes. NF- Y can function as both an activator and a repressor, depending on its interacting cofactors.