

Nkx-3.1 Polyclonal Antibody
Catalog # AP71325**Specification**

Nkx-3.1 Polyclonal Antibody - Product Information

Application	WB, IHC-P
Primary Accession	Q99801
Reactivity	Human
Host	Rabbit
Clonality	Polyclonal

Nkx-3.1 Polyclonal Antibody - Additional Information**Gene ID** 4824**Other Names**

NKX3-1; NKX3.1; NKX3A; Homeobox protein Nkx-3.1; Homeobox protein NK-3 homolog A

Dilution

WB~~Western Blot: 1/500 - 1/2000. Immunohistochemistry: 1/100 - 1/300. ELISA: 1/20000. Not yet tested in other applications.

IHC-P~~N/A

Format

Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.09% (W/V) sodium azide.

Storage Conditions

-20°C

Nkx-3.1 Polyclonal Antibody - Protein Information**Name** NKX3-1 ([HGNC:7838](#))**Function**

Transcription factor, which binds preferentially the consensus sequence 5'-TAAGT[AG]-3' and can behave as a transcriptional repressor. Plays an important role in normal prostate development, regulating proliferation of glandular epithelium and in the formation of ducts in prostate. Acts as a tumor suppressor controlling prostate carcinogenesis, as shown by the ability to inhibit proliferation and invasion activities of PC-3 prostate cancer cells.

Cellular Location

Nucleus {ECO:0000255|PROSITE-ProRule:PRU00108, ECO:0000269|PubMed:11137288}

Tissue Location


Highly expressed in the prostate and, at a lower level, in the testis.

Nkx-3.1 Polyclonal Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

Nkx-3.1 Polyclonal Antibody - Images



Western Blot analysis of various cells using Nkx-3.1 Polyclonal Antibody cells nucleus extracted by Minute TM Cytoplasmic and Nuclear Fractionation kit (SC-003, Inventbiotech, MN, USA).

Nkx-3.1 Polyclonal Antibody - Background

Transcription factor, which binds preferentially the consensus sequence 5'-TAAGT[AG]-3' and can behave as a transcriptional repressor. Plays an important role in normal prostate development, regulating proliferation of glandular epithelium and in the formation of ducts in prostate. Acts as a tumor suppressor controlling prostate carcinogenesis, as shown by the ability to inhibit proliferation and invasion activities of PC-3 prostate cancer cells.