

NRBF-2 Polyclonal Antibody
Catalog # AP71370**Specification**

NRBF-2 Polyclonal Antibody - Product Information

Application	WB
Primary Accession	Q96F24
Reactivity	Human, Mouse, Rat
Host	Rabbit
Clonality	Polyclonal

NRBF-2 Polyclonal Antibody - Additional Information**Gene ID** 29982**Other Names**

NRBF2; COPR; Nuclear receptor-binding factor 2; NRBF-2; Comodulator of PPAR and RXR

Dilution

WB~~Western Blot: 1/500 - 1/2000. ELISA: 1/20000. Not yet tested in other applications.

Format

Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.09% (W/V) sodium azide.

Storage Conditions

-20°C

NRBF-2 Polyclonal Antibody - Protein Information**Name** NRBF2**Synonyms** COPR**Function**

May modulate transcriptional activation by target nuclear receptors. Can act as transcriptional activator (in vitro).

Cellular Location

Nucleus {ECO:0000250|UniProtKB:Q9QYK3}. Cytoplasm {ECO:0000250|UniProtKB:Q9QYK3}. Cytoplasmic vesicle. Cytoplasmic vesicle, autophagosome

Tissue Location



Detected in keratinocytes, liver and placenta (PubMed:15610520). Expressed in a subset of cells in pediatric medulloblastoma (PubMed:18619852).

NRBF-2 Polyclonal Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

NRBF-2 Polyclonal Antibody - Images



NRBF-2 Polyclonal Antibody - Background

May modulate transcriptional activation by target nuclear receptors. Can act as transcriptional activator (in vitro).