

### **NXF3 Polyclonal Antibody**

**Catalog # AP71397** 

# **Specification**

### **NXF3 Polyclonal Antibody - Product Information**

Application WB, IHC-P
Primary Accession
Reactivity Human
Host Rabbit
Clonality Polyclonal

# **NXF3 Polyclonal Antibody - Additional Information**

**Gene ID 56000** 

**Other Names** 

NXF3; TAPL3; Nuclear RNA export factor 3; TAP-like protein 3; TAPL-3

**Dilution** 

WB $\sim\sim$ Western Blot: 1/500 - 1/2000. Immunohistochemistry: 1/100 - 1/300. ELISA: 1/40000. Not yet tested in other applications.

IHC-P~~N/A

#### **Format**

Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.09% (W/V) sodium azide.

## **Storage Conditions**

-20°C

#### **NXF3 Polyclonal Antibody - Protein Information**

Name NXF3

**Synonyms TAPL3** 

#### **Function**

May function as a tissue-specific nuclear mRNA export factor.


### **Cellular Location**

Nucleus. Cytoplasm. Note=Shuttles between the nucleus and the cytoplasm

### **Tissue Location**

Expressed at high level in testis and at low level in a small number of tissues

#### **NXF3 Polyclonal Antibody - Protocols**






Provided below are standard protocols that you may find useful for product applications.

- Western Blot
- Blocking Peptides
- Dot Blot
- <u>Immunohistochemistry</u>
- <u>Immunofluorescence</u>
- <u>Immunoprecipitation</u>
- Flow Cytomety
- Cell Culture

## **NXF3 Polyclonal Antibody - Images**



Immunohistochemical analysis of paraffin-embedded Human brain. Antibody was diluted at 1:100(4°,overnight). High-pressure and temperature Tris-EDTA,pH8.0 was used for antigen retrieval. Negetive contrl (right) obtaned from antibody was pre-absorbed by immunogen peptide.

## **NXF3 Polyclonal Antibody - Background**

May function as a tissue-specific nuclear mRNA export factor.