

LMTK2 Antibody (Center)
Purified Rabbit Polyclonal Antibody (Pab)
Catalog # AP7140c

Specification

LMTK2 Antibody (Center) - Product Information

Application	WB,E
Primary Accession	Q8IWU2
Reactivity	Human
Host	Rabbit
Clonality	Polyclonal
Isotype	Rabbit IgG
Calculated MW	164900
Antigen Region	668-697

LMTK2 Antibody (Center) - Additional Information

Gene ID 22853

Other Names

Serine/threonine-protein kinase LMTK2, Apoptosis-associated tyrosine kinase 2, Brain-enriched kinase, hBREK, CDK5/p35-regulated kinase, CPRK, Kinase/phosphatase/inhibitor 2, Lemur tyrosine kinase 2, Serine/threonine-protein kinase KPI-2, LMTK2, AATYK2, BREK, KIAA1079, KPI2, LMR2

Target/Specificity

This LMTK2 antibody is generated from rabbits immunized with a KLH conjugated synthetic peptide between 668-697 amino acids from the Central region of human LMTK2.

Dilution

WB~~1:1000

Format

Purified polyclonal antibody supplied in PBS with 0.09% (W/V) sodium azide. This antibody is prepared by Saturated Ammonium Sulfate (SAS) precipitation followed by dialysis against PBS.

Storage

Maintain refrigerated at 2-8°C for up to 2 weeks. For long term storage store at -20°C in small aliquots to prevent freeze-thaw cycles.

Precautions

LMTK2 Antibody (Center) is for research use only and not for use in diagnostic or therapeutic procedures.

LMTK2 Antibody (Center) - Protein Information

Name LMTK2

Synonyms AATYK2, BREK, KIAA1079, KPI2, LMR2

Function Phosphorylates PPP1C, phosphorylase b and CFTR.

Cellular Location

Membrane; Multi- pass membrane protein

Tissue Location


Mainly expressed in skeletal muscle, and weakly in brain and pancreas.

LMTK2 Antibody (Center) - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

LMTK2 Antibody (Center) - Images



Western blot analysis of LMTK2 (arrow) using rabbit polyclonal LMTK2 Antibody (Center) (Cat.#AP7140c).293 cell lysates (2 ug/lane) either nontransfected (Lane 1) or transiently transfected with the LMTK2 gene (Lane 2) (Origene Technologies).

LMTK2 Antibody (Center) - Background

Protein kinases are enzymes that transfer a phosphate group from a phosphate donor, generally the γ phosphate of ATP, onto an acceptor amino acid in a substrate protein. By this basic mechanism, protein kinases mediate most of the signal transduction in eukaryotic cells, regulating cellular metabolism, transcription, cell cycle progression, cytoskeletal rearrangement and cell movement, apoptosis, and differentiation. With more than 500 gene products, the protein kinase family is one of the largest families of proteins in eukaryotes. The family has been classified in 8 major groups based on sequence comparison of their tyrosine (PTK) or serine/threonine (STK) kinase catalytic domains. The tyrosine kinase (TK) group is mainly involved in the regulation of cell-cell interactions such as differentiation, adhesion, motility and death. There are currently about 90 TK genes sequenced, 58 are of receptor protein TK (e.g. EGFR, EPH, FGFR, PDGFR, TRK, and VEGFR families), and 32 of cytosolic TK (e.g. ABL, FAK, JAK, and SRC families).

LMTK2 Antibody (Center) - References

- Kawa, S., et al., Genes Cells 9(3):219-232 (2004).
Hillier, L.W., et al., Nature 424(6945):157-164 (2003).
Scherer, S.W., et al., Science 300(5620):767-772 (2003).
Kesavapany, S., et al., J. Neurosci. 23(12):4975-4983 (2003).
Wang, H., et al., J. Biol. Chem. 277(51):49605-49612 (2002).