

Oct-1 Polyclonal Antibody
Catalog # AP71413**Specification**

Oct-1 Polyclonal Antibody - Product Information

Application	WB
Primary Accession	P14859
Reactivity	Human, Mouse
Host	Rabbit
Clonality	Polyclonal

Oct-1 Polyclonal Antibody - Additional Information**Gene ID** 5451**Other Names**

POU2F1; OCT1; OTF1; POU domain; class 2, transcription factor 1; NF-A1; Octamer-binding protein 1; Oct-1; Octamer-binding transcription factor 1; OTF-1

Dilution

WB~~Western Blot: 1/500 - 1/2000. Immunohistochemistry: 1/100 - 1/300. ELISA: 1/5000. Not yet tested in other applications.

Format

Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.09% (W/V) sodium azide.

Storage Conditions

-20°C

Oct-1 Polyclonal Antibody - Protein Information**Name** POU2F1**Synonyms** OCT1, OTF1**Function**

Transcription factor that binds to the octamer motif (5'- ATTTGCAT-3') and activates the promoters of the genes for some small nuclear RNAs (snRNA) and of genes such as those for histone H2B and immunoglobulins. Modulates transcription transactivation by NR3C1, AR and PGR.

Cellular Location

Nucleus.

Tissue Location



Ubiquitous. Isoform 2 is lymphocyte-specific.



Oct-1 Polyclonal Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

Oct-1 Polyclonal Antibody - Images





Oct-1 Polyclonal Antibody - Background

Transcription factor that binds to the octamer motif (5'-ATTGTCAT-3') and activates the promoters of the genes for some small nuclear RNAs (snRNA) and of genes such as those for histone H2B and immunoglobulins. Modulates transcription transactivation by NR3C1, AR and PGR (By similarity). In case of human herpes simplex virus (HSV) infection, POU2F1 forms a multiprotein-DNA complex with the viral transactivator protein VP16 and HCFC1 thereby enabling the transcription of the viral immediate early genes.