

Oct-2 Polyclonal Antibody
Catalog # AP71414**Specification**

Oct-2 Polyclonal Antibody - Product Information

Application	WB, IHC-P, IF
Primary Accession	P09086
Reactivity	Human, Mouse, Monkey
Host	Rabbit
Clonality	Polyclonal

Oct-2 Polyclonal Antibody - Additional Information**Gene ID** 5452**Other Names**

POU2F2; OCT2; OTF2; POU domain; class 2, transcription factor 2; Lymphoid-restricted immunoglobulin octamer-binding protein NF-A2; Octamer-binding protein 2; Oct-2; Octamer-binding transcription factor 2; OTF-2

Dilution

WB~~Western Blot: 1/500 - 1/2000. Immunohistochemistry: 1/100 - 1/300. Immunofluorescence: 1/200 - 1/1000. ELISA: 1/20000. Not yet tested in other applications.
IHC-P~~N/A
IF~~1:50~200

Format

Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.09% (W/V) sodium azide.

Storage Conditions

-20°C

Oct-2 Polyclonal Antibody - Protein Information**Name** POU2F2 ([HGNC:9213](#))**Synonyms** OCT2, OTF2**Function**

Transcription factor that specifically binds to the octamer motif (5'-ATTTGCAT-3') (PubMed:2904654, PubMed:7859290). Regulates IL6 expression in B cells with POU2AF1 (By similarity). Regulates transcription in a number of tissues in addition to activating immunoglobulin gene expression (PubMed:2901913, PubMed:2904654). Modulates transcription transactivation by NR3C1, AR and PGR (PubMed:10480874).

Cellular Location

Cytoplasm. Nucleus.


Tissue Location

Isoform 3 is B-cell specific. Isoform 5 is expressed in B-cells and the immunoglobulin-expressing T-cell line MOLT-4, but not in the T-cell line BW5147.

Oct-2 Polyclonal Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

Oct-2 Polyclonal Antibody - Images

Western Blot analysis of various cells using Oct-2 Polyclonal Antibody

Oct-2 Polyclonal Antibody - Background

Transcription factor that specifically binds to the octamer motif (5'-ATTGTCAT-3'). Regulates transcription in a number of tissues in addition to activating immunoglobulin gene expression. Modulates transcription transactivation by NR3C1, AR and PGR. Isoform 5 activates the U2 small nuclear RNA (snRNA) promoter.