

p16 Polyclonal Antibody

Catalog # AP71676

Specification

p16 Polyclonal Antibody - Product Information

Application WB
Primary Accession P42771

Reactivity Human, Mouse Host Rabbit

p16 Polyclonal Antibody - Additional Information

Gene ID 1029

Other Names

CDKN2A; CDKN2; MTS1; Cyclin-dependent kinase inhibitor 2A; isoforms 1/2/3; Cyclin-dependent kinase 4 inhibitor A; CDK4I; Multiple tumor suppressor 1; MTS-1; p16-INK4a; p16-INK4; p16INK4A

Polyclonal

Dilution

Clonality

WB~~Western Blot: 1/500 - 1/2000. ELISA: 1/20000. Not yet tested in other applications.

Format

Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.09% (W/V) sodium azide.

Storage Conditions

-20°C

p16 Polyclonal Antibody - Protein Information

Name CDKN2A (HGNC:1787)

Synonyms CDKN2, MTS1

Function

Acts as a negative regulator of the proliferation of normal cells by interacting strongly with CDK4 and CDK6. This inhibits their ability to interact with cyclins D and to phosphorylate the retinoblastoma protein.


Cellular Location


Cytoplasm. Nucleus

Tissue Location

Widely expressed but not detected in brain or skeletal muscle. Isoform 3 is pancreas-specific

p16 Polyclonal Antibody - Protocols








Provided below are standard protocols that you may find useful for product applications.

- Western Blot
- Blocking Peptides
- Dot Blot
- <u>Immunohistochemistry</u>
- Immunofluorescence
- Immunoprecipitation
- Flow Cytomety
- Cell Culture

p16 Polyclonal Antibody - Images



Western Blot analysis of various cells using p16 Polyclonal Antibody

p16 Polyclonal Antibody - Background

Acts as a negative regulator of the proliferation of normal cells by interacting strongly with CDK4 and CDK6. This inhibits their ability to interact with cyclins D and to phosphorylate the retinoblastoma protein.