

Patched Polyclonal Antibody
Catalog # AP71781**Specification**

Patched Polyclonal Antibody - Product Information

Application	WB, IHC-P
Primary Accession	Q13635
Reactivity	Human, Mouse
Host	Rabbit
Clonality	Polyclonal

Patched Polyclonal Antibody - Additional Information**Gene ID** 5727**Other Names**

PTCH1; PTCH; Protein patched homolog 1; PTC; PTC1

Dilution

WB~~Western Blot: 1/500 - 1/2000. Immunohistochemistry: 1/100 - 1/300. ELISA: 1/10000. Not yet tested in other applications.

IHC-P~~N/A

Format

Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.09% (W/V) sodium azide.

Storage Conditions

-20°C

Patched Polyclonal Antibody - Protein Information**Name** PTCH1**Synonyms** PTCH**Function**

Acts as a receptor for sonic hedgehog (SHH), indian hedgehog (IHH) and desert hedgehog (DHH). Associates with the smoothened protein (SMO) to transduce the hedgehog's proteins signal. Seems to have a tumor suppressor function, as inactivation of this protein is probably a necessary, if not sufficient step for tumorigenesis.

Cellular Location

Cell membrane {ECO:0000250|UniProtKB:Q61115}; Multi-pass membrane protein

Tissue Location



In the adult, expressed in brain, lung, liver, heart, placenta, skeletal muscle, pancreas and kidney. Expressed in tumor cells but not in normal skin




Patched Polyclonal Antibody - Protocols




Provided below are standard protocols that you may find useful for product applications.




- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)


Patched Polyclonal Antibody - Images











Patched Polyclonal Antibody - Background

Acts as a receptor for sonic hedgehog (SHH), indian hedgehog (IHH) and desert hedgehog (DHH). Associates with the smoothened protein (SMO) to transduce the hedgehog's proteins signal. Seems to have a tumor suppressor function, as inactivation of this protein is probably a necessary, if not sufficient step for tumorigenesis.