

PWWP2B Polyclonal Antibody
Catalog # AP72092**Specification**

PWWP2B Polyclonal Antibody - Product Information

Application	WB, IHC-P
Primary Accession	Q6NUJ5
Reactivity	Human
Host	Rabbit
Clonality	Polyclonal

PWWP2B Polyclonal Antibody - Additional Information**Gene ID** 170394**Other Names**

PWWP2B; PWWP2; PWWP domain-containing protein 2B

Dilution

WB~~Western Blot: 1/500 - 1/2000. Immunohistochemistry: 1/100 - 1/300. ELISA: 1/20000. Not yet tested in other applications.

IHC-P~~N/A

Format

Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.09% (W/V) sodium azide.

Storage Conditions

-20°C

PWWP2B Polyclonal Antibody - Protein Information**Name** PWWP2B**Synonyms** PWWP2**Function**

Chromatin-binding protein that acts as an adapter between distinct nucleosome components (H3K36me3 or H2A.Z) and chromatin- modifying complexes, contributing to the regulation of the levels of histone acetylation at actively transcribed genes (PubMed:30228260). Competes with CHD4 and MBD3 for interaction with MTA1 to form a NuRD subcomplex, preventing the formation of full NuRD complex (containing CHD4 and MBD3), leading to recruitment of HDACs to gene promoters resulting in turn in the deacetylation of nearby H3K27 and H2A.Z (PubMed:30228260). Plays a role in facilitating transcriptional elongation through regulation of histone acetylation (By similarity). Negatively regulates brown adipocyte thermogenesis by interacting with and stabilizing HDAC1 at the UCP1 gene promoter, thereby promoting histone deacetylation at the promoter leading to the repression of UCP1 expression (By similarity).

PWWP2B Polyclonal Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

PWWP2B Polyclonal Antibody - Images

