Relaxin Receptor 2 Polyclonal Antibody
Catalog # AP72225

Specification

Relaxin Receptor 2 Polyclonal Antibody - Product Information

Application	**WB, IF**
Primary Accession	**Q8WXD0**
Reactivity	**Human, Mouse, Rat**
Host	**Rabbit**
Clonality	**Polyclonal**

Relaxin Receptor 2 Polyclonal Antibody - Additional Information

Gene ID 122042

Other Names
RXFP2; GPR106; GREAT; LGR8; Relaxin receptor 2; G-protein coupled receptor 106; G-protein coupled receptor affecting testicular descent; Leucine-rich repeat-containing G-protein coupled receptor 8; Relaxin family peptide receptor 2

Dilution
WB~~Western Blot: 1/500 - 1/2000. Immunofluorescence: 1/200 - 1/1000. ELISA: 1/10000. Not yet tested in other applications.
IF~~1:50~200

Format
Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.09% (W/V) sodium azide.

Storage Conditions
-20℃

Relaxin Receptor 2 Polyclonal Antibody - Protein Information

Name RXFP2

Synonyms GPR106, GREAT, LGR8

Function
Receptor for relaxin. The activity of this receptor is mediated by G proteins leading to stimulation of adenylate cyclase and an increase of cAMP. May also be a receptor for Leydig insulin-like peptide (INSL3).

Cellular Location
Cell membrane; Multi-pass membrane protein.

Tissue Location
Expressed mainly in the brain, kidney, muscle, testis, thyroid, uterus, peripheral blood cells and bone marrow

Relaxin Receptor 2 Polyclonal Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- Western Blot
- Blocking Peptides
- Dot Blot
- Immunohistochemistry
- Immunofluorescence
- Immunoprecipitation
- Flow Cytomety
- Cell Culture

Relaxin Receptor 2 Polyclonal Antibody - Images

Relaxin Receptor 2 Polyclonal Antibody - Background

Receptor for relaxin. The activity of this receptor is mediated by G proteins leading to stimulation of adenylate cyclase and an increase of cAMP. May also be a receptor for Leydig insulin-like peptide (INSL3).