

# RhoGEF p115 Polyclonal Antibody

Catalog # AP72271

# **Specification**

# **RhoGEF p115 Polyclonal Antibody - Product Information**

Application WB
Primary Accession O92888

Reactivity Human, Mouse, Rat

Host Rabbit Clonality Polyclonal

# RhoGEF p115 Polyclonal Antibody - Additional Information

### **Gene ID 9138**

### **Other Names**

ARHGEF1; Rho guanine nucleotide exchange factor 1; 115 kDa guanine nucleotide exchange factor; p115-RhoGEF; p115RhoGEF; Sub1.5

#### Dilution

WB~~Western Blot: 1/500 - 1/2000. ELISA: 1/40000. Not yet tested in other applications.

### **Format**

Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.09% (W/V) sodium azide.

# **Storage Conditions**

-20°C

### RhoGEF p115 Polyclonal Antibody - Protein Information

### Name ARHGEF1


# **Function**

Seems to play a role in the regulation of RhoA GTPase by guanine nucleotide-binding alpha-12 (GNA12) and alpha-13 (GNA13) subunits (PubMed:<a

href="http://www.uniprot.org/citations/9641915" target="\_blank">9641915</a>, PubMed:<a href="http://www.uniprot.org/citations/9641916" target="\_blank">9641916</a>). Acts as a GTPase-activating protein (GAP) for GNA12 and GNA13, and as guanine nucleotide exchange factor (GEF) for RhoA GTPase (PubMed:<a href="http://www.uniprot.org/citations/30521495" target="\_blank">30521495</a>, PubMed:<a href="http://www.uniprot.org/citations/8810315" target="\_blank">8810315</a>, PubMed:<a href="http://www.uniprot.org/citations/9641915" target="\_blank">9641915</a>, PubMed:<a href="http://www.uniprot.org/citations/9641916" target="\_blank">9641916</a>, PubMed:<a href="http://www.uniprot.org/citations/9641916" target="\_blank">9641916</a>). Activated G alpha 13/GNA13 stimulates the RhoGEF activity through interaction with the RGS-like domain (PubMed:<a

href="http://www.uniprot.org/citations/9641916" target="\_blank">9641916</a>). This GEF activity is inhibited by binding to activated GNA12 (PubMed:<a

href="http://www.uniprot.org/citations/9641916" target="\_blank">9641916</a>). Mediates angiotensin-2-induced RhoA activation (PubMed:<a





href="http://www.uniprot.org/citations/20098430" target="\_blank">20098430</a>). In lymphoid follicles, may trigger activation of GNA13 as part of S1PR2-dependent signaling pathway that leads to inhibition of germinal center (GC) B cell growth and migration outside the GC niche.

### **Cellular Location**

Cytoplasm. Membrane. Note=Translocated to the membrane by activated GNA13 or LPA stimulation

### **Tissue Location**


Ubiquitously expressed.

# **RhoGEF p115 Polyclonal Antibody - Protocols**


Provided below are standard protocols that you may find useful for product applications.

- Western Blot
- Blocking Peptides
- Dot Blot
- <u>Immunohistochemistry</u>
- Immunofluorescence
- Immunoprecipitation
- Flow Cytomety
- Cell Culture

# **RhoGEF p115 Polyclonal Antibody - Images**







**RhoGEF p115 Polyclonal Antibody - Background** 

Seems to play a role in the regulation of RhoA GTPase by guanine nucleotide-binding alpha-12 (GNA12) and alpha-13 (GNA13) subunits. Acts as GTPase-activating protein (GAP) for GNA12 and GNA13, and as guanine nucleotide exchange factor (GEF) for RhoA GTPase. Activated G alpha 13/GNA13 stimulates the RhoGEF activity through interaction with the RGS-like domain. This GEF activity is inhibited by binding to activated GNA12. Mediates angiotensin-2- induced RhoA activation.