

**SAP 14 Polyclonal Antibody**  
**Catalog # AP72393****Specification**

---

**SAP 14 Polyclonal Antibody - Product Information**

Application	WB
Primary Accession	<a href="#">Q9Y3B4</a>
Reactivity	Human, Mouse
Host	Rabbit
Clonality	Polyclonal

**SAP 14 Polyclonal Antibody - Additional Information****Gene ID** 51639**Other Names**

SF3B14; CGI-110; HSPC175; HT006; Pre-mRNA branch site protein p14; SF3b 14 kDa subunit

**Dilution**

WB~~Western Blot: 1/500 - 1/2000. ELISA: 1/40000. Not yet tested in other applications.

**Format**

Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.09% (W/V) sodium azide.

**Storage Conditions**

-20°C

**SAP 14 Polyclonal Antibody - Protein Information****Name** SF3B6**Synonyms** SAP14, SF3B14, SF3B14A**Function**

Component of the 17S U2 SnRNP complex of the spliceosome, a large ribonucleoprotein complex that removes introns from transcribed pre-mRNAs (PubMed:<a href="http://www.uniprot.org/citations/27720643" target="\_blank">27720643</a>, PubMed:<a href="http://www.uniprot.org/citations/12234937" target="\_blank">12234937</a>, PubMed:<a href="http://www.uniprot.org/citations/32494006" target="\_blank">32494006</a>, PubMed:<a href="http://www.uniprot.org/citations/34822310" target="\_blank">34822310</a>). The 17S U2 SnRNP complex (1) directly participates in early spliceosome assembly and (2) mediates recognition of the intron branch site during pre-mRNA splicing by promoting the selection of the pre-mRNA branch-site adenosine, the nucleophile for the first step of splicing (PubMed:<a href="http://www.uniprot.org/citations/12234937" target="\_blank">12234937</a>, PubMed:<a href="http://www.uniprot.org/citations/32494006" target="\_blank">32494006</a>, PubMed:<a href="http://www.uniprot.org/citations/34822310" target="\_blank">34822310</a>). Within the 17S U2 SnRNP complex, SF3B6 is part of the SF3B subcomplex, which is required for 'A' complex assembly formed by the stable binding of U2 snRNP to the branchpoint sequence in pre-mRNA

(PubMed:<a href="http://www.uniprot.org/citations/12234937" target="\_blank">12234937</a>, PubMed:<a href="http://www.uniprot.org/citations/27720643" target="\_blank">27720643</a>). Sequence independent binding of SF3A and SF3B subcomplexes upstream of the branch site is essential, it may anchor U2 snRNP to the pre-mRNA (PubMed:<a href="http://www.uniprot.org/citations/12234937" target="\_blank">12234937</a>). Within the 17S U2 SnRNP complex, SF3B6 directly contacts the pre-mRNA branch site adenosine for the first catalytic step of splicing (PubMed:<a href="http://www.uniprot.org/citations/16432215" target="\_blank">16432215</a>). SF3B6 stabilizes the intron branch site-U2 snRNA duplex, thereby promoting- binding of introns with poor sequence complementarity (PubMed:<a href="http://www.uniprot.org/citations/34822310" target="\_blank">34822310</a>). Also acts as a component of the minor spliceosome, which is involved in the splicing of U12-type introns in pre-mRNAs (PubMed:<a href="http://www.uniprot.org/citations/15146077" target="\_blank">15146077</a>, PubMed:<a href="http://www.uniprot.org/citations/33509932" target="\_blank">33509932</a>).

### Cellular Location


Nucleus


### SAP 14 Polyclonal Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

### SAP 14 Polyclonal Antibody - Images





#### **SAP 14 Polyclonal Antibody - Background**

Involved in pre-mRNA splicing as a component of the splicing factor SF3B complex (PubMed:27720643). SF3B complex is required for 'A' complex assembly formed by the stable binding of U2 snRNP to the branchpoint sequence (BPS) in pre-mRNA (PubMed:12234937). Directly contacts the pre-mRNA branch site adenosine for the first catalytic step of splicing (PubMed:16432215). Enters the spliceosome and associates with the pre-mRNA branch site as part of the 17S U2 or, in the case of the minor spliceosome, as part of the 18S U11/U12 snRNP complex, and thus may facilitate the interaction of these snRNP with the branch sites of U2 and U12 respectively (PubMed:16432215).