

SR-1A Polyclonal Antibody

Catalog # AP72574

Specification

SR-1A Polyclonal Antibody - Product Information

Application WB, IHC-P, IF
Primary Accession
Reactivity Human, Mouse, Rat
Host Rabbit
Clonality Polyclonal

SR-1A Polyclonal Antibody - Additional Information

Gene ID 3350

Other Names

HTR1A; ADRB2RL1; ADRBRL1; 5-hydroxytryptamine receptor 1A; 5-HT-1A; 5-HT1A; G-21; Serotonin receptor 1A

Dilution

WB~~Western Blot: 1/500 - 1/2000. Immunohistochemistry: 1/100 - 1/300. Immunofluorescence: 1/200 - 1/1000. ELISA: 1/20000. Not yet tested in other applications. IHC-P~~N/A IF~~1:50~200

Format

Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.09% (W/V) sodium azide.

Storage Conditions

-20°C

SR-1A Polyclonal Antibody - Protein Information

Name HTR1A (HGNC:5286)

Synonyms ADRB2RL1, ADRBRL1

Function

G-protein coupled receptor for 5-hydroxytryptamine (serotonin) (PubMed:22957663, PubMed:3138543, PubMed:33762731, PubMed:37935376, PubMed:37935377, PubMed:8138923, PubMed:8393041). Also functions as a receptor for various drugs and psychoactive substances (PubMed:22957663, PubMed:22957663, PubMed:<a



href="http://www.uniprot.org/citations/3138543" target=" blank">3138543, PubMed:33762731, PubMed:38552625, PubMed:8138923, PubMed:8393041). Ligand binding causes a conformation change that triggers signaling via guanine nucleotide-binding proteins (G proteins) and modulates the activity of downstream effectors, such as adenylate cyclase (PubMed:22957663, PubMed:3138543, PubMed:33762731, PubMed: 8138923, PubMed: 8393041). HTR1A is coupled to G(i)/G(o) G alpha proteins and mediates inhibitory neurotransmission: signaling inhibits adenylate cyclase activity and activates a phosphatidylinositol-calcium second messenger system that regulates the release of Ca(2+) ions from intracellular stores (PubMed:33762731, PubMed:35610220). Beta-arrestin family members regulate signaling by mediating both receptor desensitization and resensitization processes (PubMed: 18476671, PubMed:20363322, PubMed:20945968). Plays a role in the regulation of 5- hydroxytryptamine release and in the regulation of dopamine and 5- hydroxytryptamine metabolism (PubMed: 18476671, PubMed:20363322, PubMed:20945968). Plays a role in the regulation of dopamine and 5- hydroxytryptamine levels in the brain, and thereby affects neural activity, mood and behavior (PubMed: 18476671, PubMed:20363322, PubMed:20945968). Plays a role in the response to anxiogenic stimuli (PubMed: 18476671, PubMed:20363322, PubMed:20945968).

Cellular Location

Cell membrane; Multi-pass membrane protein. Cell projection, dendrite {ECO:0000250|UniProtKB:P19327}

Tissue Location

Detected in lymph nodes, thymus and spleen. Detected in activated T-cells, but not in resting T-cells


SR-1A Polyclonal Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- Western Blot
- Blocking Peptides
- Dot Blot
- <u>Immunohistochemistry</u>
- Immunofluorescence
- Immunoprecipitation
- Flow Cytomety
- Cell Culture



SR-1A Polyclonal Antibody - Images



SR-1A Polyclonal Antibody - Background

G-protein coupled receptor for 5-hydroxytryptamine (serotonin). Also functions as a receptor for various drugs and psychoactive substances. Ligand binding causes a conformation change that triggers signaling via guanine nucleotide-binding proteins (G proteins) and modulates the activity of down-stream effectors, such as adenylate cyclase. Beta-arrestin family members inhibit signaling via G proteins and mediate activation of alternative signaling pathways. Signaling inhibits adenylate cyclase activity and activates a phosphatidylinositol-calcium second messenger system that regulates the release of Ca(2+) ions from intracellular stores. Plays a role in the regulation of 5-hydroxytryptamine release and in the regulation of dopamine and 5-hydroxytryptamine levels in the brain, and thereby affects neural activity, mood and behavior. Plays a role in the response to anxiogenic stimuli.