

SREBP-1 Polyclonal Antibody
Catalog # AP72594**Specification**

SREBP-1 Polyclonal Antibody - Product Information

Application	WB, IHC-P
Primary Accession	P36956
Reactivity	Human, Rat
Host	Rabbit
Clonality	Polyclonal

SREBP-1 Polyclonal Antibody - Additional Information**Gene ID** 6720**Other Names**

SREBF1; BHLHD1; SREBP1; Sterol regulatory element-binding protein 1; SREBP-1; Class D basic helix-loop-helix protein 1; bHLHD1; Sterol regulatory element-binding transcription factor 1

Dilution

WB~~Western Blot: 1/500 - 1/2000. Immunohistochemistry: 1/100 - 1/300. ELISA: 1/20000. Not yet tested in other applications.

IHC-P~~N/A

Format

Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.09% (W/V) sodium azide.

Storage Conditions

-20°C

SREBP-1 Polyclonal Antibody - Protein Information

Name SREBF1 {ECO:0000303|PubMed:7759101, ECO:0000312|HGNC:HGNC:11289}

Function

[Sterol regulatory element-binding protein 1]: Precursor of the transcription factor form (Processed sterol regulatory element-binding protein 1), which is embedded in the endoplasmic reticulum membrane (PubMed:<[a href="http://www.uniprot.org/citations/32322062"](http://www.uniprot.org/citations/32322062)target="_blank">32322062). Low sterol concentrations promote processing of this form, releasing the transcription factor form that translocates into the nucleus and activates transcription of genes involved in cholesterol biosynthesis and lipid homeostasis (By similarity).

Cellular Location

[Sterol regulatory element-binding protein 1]: Endoplasmic reticulum membrane; Multi-pass membrane protein. Golgi apparatus membrane; Multi-pass membrane protein. Cytoplasmic vesicle, COPII-coated vesicle membrane {ECO:0000250|UniProtKB:Q9WTN3}; Multi-pass membrane protein. Note=At high sterol concentrations, the SCAP-SREBP is retained in the endoplasmic reticulum. Low sterol concentrations promote recruitment into COPII-coated vesicles

and transport of the SCAP-SREBP to the Golgi, where it is processed
{ECO:0000250|UniProtKB:Q9WTN3} [Isoform SREBP-1aDelta]: Nucleus

Tissue Location


Expressed in a wide variety of tissues, most abundant in liver and adrenal gland (PubMed:8402897). In fetal tissues lung and liver shows highest expression (PubMed:8402897) [Isoform SREBP-1C]: Predominantly expressed in liver and adipose tissues (PubMed:8402897). Also expressed in kidney, brain, white fat, and muscle (PubMed:8402897)

SREBP-1 Polyclonal Antibody - Protocols


Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

SREBP-1 Polyclonal Antibody - Images



Western Blot analysis of various cells using SREBP-1 Polyclonal Antibody diluted at 1:1000



Western blot analysis of 3T3 lysis using SREBP-1 antibody. Antibody was diluted at 1:1000

SREBP-1 Polyclonal Antibody - Background

Transcriptional activator required for lipid homeostasis. Regulates transcription of the LDL receptor gene as well as the fatty acid and to a lesser degree the cholesterol synthesis pathway (By similarity). Binds to the sterol regulatory element 1 (SRE-1) (5'-ATCACCCCAC-3'). Has dual sequence specificity binding to both an E-box motif (5'-ATCACGTGA-3') and to SRE-1 (5'-ATCACCCCAC-3').