

SRE-ZBP Polyclonal Antibody
Catalog # AP72596**Specification**

SRE-ZBP Polyclonal Antibody - Product Information

Application	WB, IF
Primary Accession	Q16670
Reactivity	Human
Host	Rabbit
Clonality	Polyclonal

SRE-ZBP Polyclonal Antibody - Additional Information**Gene ID** 7741**Other Names**

ZNF187; ZSCAN26; Zinc finger protein 187; Protein SRE-ZBP; Zinc finger and SCAN domain-containing protein 26

Dilution

WB~~Western Blot: 1/500 - 1/2000. Immunofluorescence: 1/200 - 1/1000. ELISA: 1/5000. Not yet tested in other applications.

IF~~1:50~200

Format

Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.09% (W/V) sodium azide.

Storage Conditions

-20°C

SRE-ZBP Polyclonal Antibody - Protein Information**Name** ZSCAN26**Synonyms** ZNF187**Function**

May be involved in transcriptional regulation.

Cellular Location


Nucleus {ECO:0000255|PROSITE-ProRule:PRU00187, ECO:0000269|PubMed:1569959}

SRE-ZBP Polyclonal Antibody - Protocols


Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)


SRE-ZBP Polyclonal Antibody - Images




Western Blot analysis of various cells using SRE-ZBP Polyclonal Antibody cells nucleus extracted by Minute TM Cytoplasmic and Nuclear Fractionation kit (SC-003, Invent biotech, MN, USA).



Immunohistochemical analysis of paraffin-embedded Human brain. Antibody was diluted at 1:100 (4°, overnight). High-pressure and temperature Tris-EDTA, pH 8.0 was used for antigen retrieval. Negative control (right) obtained from antibody was pre-absorbed by immunogen peptide.



Western Blot analysis of various cells using SRE-ZBP Polyclonal Antibody cells nucleus extracted by Minute TM Cytoplasmic and Nuclear Fractionation kit (SC-003, Inventbiotech, MN, USA).



Immunohistochemical analysis of paraffin-embedded Human brain. Antibody was diluted at 1:100 (4°, overnight). High-pressure and temperature Tris-EDTA, pH 8.0 was used for antigen retrieval. Negative control (right) obtained from antibody was pre-absorbed by immunogen peptide.

SRE-ZBP Polyclonal Antibody - Background

May be involved in transcriptional regulation.