

TAF II p68 Polyclonal Antibody
Catalog # AP72713**Specification**

TAF II p68 Polyclonal Antibody - Product Information

Application	WB
Primary Accession	Q92804
Reactivity	Human, Mouse
Host	Rabbit
Clonality	Polyclonal

TAF II p68 Polyclonal Antibody - Additional Information**Gene ID** 8148**Other Names**

TAF15; RBP56; TAF2N; TATA-binding protein-associated factor 2N; 68 kDa TATA-binding protein-associated factor; TAF(II)68; TAFII68; RNA-binding protein 56

Dilution

WB~~Western Blot: 1/500 - 1/2000. Immunohistochemistry: 1/100 - 1/300. ELISA: 1/5000. Not yet tested in other applications.

Format

Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.09% (W/V) sodium azide.

Storage Conditions

-20°C

TAF II p68 Polyclonal Antibody - Protein Information**Name** TAF15**Synonyms** RBP56, TAF2N**Function**

RNA and ssDNA-binding protein that may play specific roles during transcription initiation at distinct promoters. Can enter the preinitiation complex together with the RNA polymerase II (Pol II).

Cellular Location

Nucleus. Cytoplasm. Note=Shuttles from the nucleus to the cytoplasm

Tissue Location



Ubiquitous. Observed in all fetal and adult tissues

TAF II p68 Polyclonal Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

TAF II p68 Polyclonal Antibody - Images



TAF II p68 Polyclonal Antibody - Background

RNA and ssDNA-binding protein that may play specific roles during transcription initiation at

distinct promoters. Can enter the preinitiation complex together with the RNA polymerase II (Pol II).