

Tra-2 α Polyclonal Antibody
Catalog # AP72896**Specification**

Tra-2 α Polyclonal Antibody - Product Information

Application	WB, IHC-P, IF
Primary Accession	Q13595
Reactivity	Human, Mouse
Host	Rabbit
Clonality	Polyclonal

Tra-2 α Polyclonal Antibody - Additional Information**Gene ID** 29896**Other Names**

TRA2A; Transformer-2 protein homolog alpha; TRA-2 alpha; TRA2-alpha; Transformer-2 protein homolog A

DilutionWB~~Western Blot: 1/500 - 1/2000. Immunohistochemistry: 1/100 - 1/300. Immunofluorescence: 1/200 - 1/1000. ELISA: 1/20000. Not yet tested in other applications.
IHC-P~~N/A
IF~~1:50~200**Format**

Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.09% (W/V) sodium azide.

Storage Conditions

-20°C

Tra-2 α Polyclonal Antibody - Protein Information**Name** TRA2A ([HGNC:16645](#))**Function**

Sequence-specific RNA-binding protein which participates in the control of pre-mRNA splicing.

Cellular Location

Nucleus.


Tra-2 α Polyclonal Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)

- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

Tra-2 α Polyclonal Antibody - Images



Western Blot analysis of various cells using Tra-2 α Polyclonal Antibody diluted at 1:500. Secondary antibody was diluted at 1:20000 cells nucleus extracted by Minute TM Cytoplasmic and Nuclear Fractionation kit (SC-003, Invent biotech, MN, USA).

Tra-2 α Polyclonal Antibody - Background

Sequence-specific RNA-binding protein which participates in the control of pre-mRNA splicing.