

Translin Polyclonal Antibody
Catalog # AP72905**Specification**

Translin Polyclonal Antibody - Product Information

Application	WB, IHC-P
Primary Accession	Q15631
Reactivity	Human, Mouse, Rat
Host	Rabbit
Clonality	Polyclonal

Translin Polyclonal Antibody - Additional Information**Gene ID** 7247**Other Names**

TSN; Translin; Component 3 of promoter of RISC; C3PO

Dilution

WB~~Western Blot: 1/500 - 1/2000. Immunohistochemistry: 1/100 - 1/300. ELISA: 1/40000. Not yet tested in other applications.

IHC-P~~N/A

Format

Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.09% (W/V) sodium azide.

Storage Conditions

-20°C

Translin Polyclonal Antibody - Protein Information**Name** TSN ([HGNC:12379](#))**Function**

DNA-binding protein that specifically recognizes consensus sequences at the breakpoint junctions in chromosomal translocations, mostly involving immunoglobulin (Ig)/T-cell receptor gene segments. Seems to recognize single-stranded DNA ends generated by staggered breaks occurring at recombination hot spots.

Cellular Location


Cytoplasm. Nucleus

Translin Polyclonal Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

Translin Polyclonal Antibody - Images



Translin Polyclonal Antibody - Background

DNA-binding protein that specifically recognizes consensus sequences at the breakpoint junctions in chromosomal translocations, mostly involving immunoglobulin (Ig)/T-cell receptor gene segments. Seems to recognize single-stranded DNA ends generated by staggered breaks occurring at recombination hot spots.