

**USP30 Polyclonal Antibody**  
**Catalog # AP73017****Specification****USP30 Polyclonal Antibody - Product Information**

|                   |                        |
|-------------------|------------------------|
| Application       | WB                     |
| Primary Accession | <a href="#">Q70CQ3</a> |
| Reactivity        | Human, Mouse, Rat      |
| Host              | Rabbit                 |
| Clonality         | Polyclonal             |

**USP30 Polyclonal Antibody - Additional Information****Gene ID** 84749**Other Names**

USP30; Ubiquitin carboxyl-terminal hydrolase 30; Deubiquitinating enzyme 30; Ubiquitin thioesterase 30; Ubiquitin-specific-processing protease 30; Ub-specific protease 30

**Dilution**

WB~~Western Blot: 1/500 - 1/2000. ELISA: 1/10000. Not yet tested in other applications.

**Format**

Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.09% (W/V) sodium azide.

**Storage Conditions**

-20°C

**USP30 Polyclonal Antibody - Protein Information****Name** USP30 ([HGNC:20065](#))**Function**

Deubiquitinating enzyme tethered to the mitochondrial outer membrane that acts as a key inhibitor of mitophagy by counteracting the action of parkin (PRKN): hydrolyzes ubiquitin attached by parkin on target proteins, such as RHOT1/MIRO1 and TOMM20, thereby blocking parkin's ability to drive mitophagy (PubMed:<a href="http://www.uniprot.org/citations/18287522" target="\_blank">18287522</a>, PubMed:<a href="http://www.uniprot.org/citations/24896179" target="\_blank">24896179</a>, PubMed:<a href="http://www.uniprot.org/citations/25527291" target="\_blank">25527291</a>, PubMed:<a href="http://www.uniprot.org/citations/25621951" target="\_blank">25621951</a>). Preferentially cleaves 'Lys-6'- and 'Lys-11'-linked polyubiquitin chains, 2 types of linkage that participate in mitophagic signaling (PubMed:<a href="http://www.uniprot.org/citations/25621951" target="\_blank">25621951</a>). Does not cleave efficiently polyubiquitin phosphorylated at 'Ser-65' (PubMed:<a href="http://www.uniprot.org/citations/25527291" target="\_blank">25527291</a>). Acts as a negative regulator of mitochondrial fusion by mediating deubiquitination of MFN1 and MFN2 (By similarity).

**Cellular Location**

Mitochondrion outer membrane


**Tissue Location**

Expressed in skeletal muscle, pancreas, liver and kidney.

**USP30 Polyclonal Antibody - Protocols**

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

**USP30 Polyclonal Antibody - Images****USP30 Polyclonal Antibody - Background**

Deubiquitinating enzyme tethered to the mitochondrial outer membrane that acts as a key inhibitor of mitophagy by counteracting the action of parkin (PRKN): hydrolyzes ubiquitin attached by parkin on target proteins, such as RHOT1/MIRO1 and TOMM20, thereby blocking parkin's ability to drive mitophagy (PubMed:18287522, PubMed:24896179, PubMed:25527291, PubMed:25621951). Preferentially cleaves 'Lys-6'- and 'Lys-11'- linked polyubiquitin chains, 2 types of linkage that participate to mitophagic signaling (PubMed:25621951). Does not cleave efficiently polyubiquitin phosphorylated at 'Ser-65' (PubMed:25527291). Acts as negative regulator of mitochondrial fusion by mediating deubiquitination of MFN1 and MFN2 (By similarity).