

WASP Polyclonal Antibody

Catalog # AP73079

Specification

WASP Polyclonal Antibody - Product Information

Application WB
Primary Accession P42768

Reactivity Human, Mouse

Host Rabbit Clonality Polyclonal

WASP Polyclonal Antibody - Additional Information

Gene ID 7454

Other Names

WAS; IMD2; Wiskott-Aldrich syndrome protein; WASp

Dilution

WB~~Western Blot: 1/500 - 1/2000. Immunohistochemistry: 1/100 - 1/300. ELISA: 1/5000. Not yet tested in other applications.

Format

Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.09% (W/V) sodium azide.

Storage Conditions

-20°C


WASP Polyclonal Antibody - Protein Information

Name WAS

Synonyms IMD2

Function

Effector protein for Rho-type GTPases that regulates actin filament reorganization via its interaction with the Arp2/3 complex (PubMed:12235133, PubMed:12769847, PubMed:16275905). Important for efficient actin polymerization (PubMed:8625410, PubMed:12235133, PubMed:16275905). Possible regulator of lymphocyte and platelet function (PubMed:9405671). Mediates actin filament reorganization and the formation of actin pedestals upon infection by pathogenic bacteria (PubMed:18650809). In addition to its role in the cytoplasmic cytoskeleton, also





Tel: 858.875.1900 Fax: 858.875.1999

promotes actin polymerization in the nucleus, thereby regulating gene transcription and repair of damaged DNA (PubMed:20574068). Promotes homologous recombination (HR) repair in response to DNA damage by promoting nuclear actin polymerization, leading to drive motility of double-strand breaks (DSBs) (PubMed: 29925947).

Cellular Location

Cytoplasm, cytoskeleton. Nucleus

Tissue Location


Expressed predominantly in the thymus. Also found, to a much lesser extent, in the spleen.

WASP Polyclonal Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- Western Blot
- Blocking Peptides
- Dot Blot
- Immunohistochemistry
- Immunofluorescence
- Immunoprecipitation
- Flow Cytomety
- Cell Culture

WASP Polyclonal Antibody - Images



WASP Polyclonal Antibody - Background

Effector protein for Rho-type GTPases that regulates actin filament reorganization via its interaction with the Arp2/3 complex (PubMed:12235133, PubMed:12769847, PubMed:16275905). Important for efficient actin polymerization (PubMed:8625410, PubMed:12235133, PubMed:16275905). Possible regulator of lymphocyte and platelet function (PubMed:9405671). Mediates actin filament reorganization and the formation of actin pedestals upon infection by





Tel: 858.875.1900 Fax: 858.875.1999

pathogenic bacteria (PubMed:18650809). In addition to its role in the cytoplasmic cytoskeleton, also promotes actin polymerization in the nucleus, thereby regulating gene transcription and repair of damaged DNA (PubMed:20574068). Promotes homologous recombination (HR) repair in response to DNA damage by promoting nuclear actin polymerization, leading to drive motility of double-strand breaks (DSBs) (PubMed:29925947).