

ZADH2 Polyclonal Antibody
Catalog # AP73111**Specification**

ZADH2 Polyclonal Antibody - Product Information

Application	WB, IHC-P
Primary Accession	Q8N4Q0
Reactivity	Human, Mouse, Rat
Host	Rabbit
Clonality	Polyclonal

ZADH2 Polyclonal Antibody - Additional Information**Gene ID** 284273**Other Names**

ZADH2; Zinc-binding alcohol dehydrogenase domain-containing protein 2

Dilution

WB~~Western Blot: 1/500 - 1/2000. Immunohistochemistry: 1/100 - 1/300. ELISA: 1/10000. Not yet tested in other applications.

IHC-P~~N/A

Format

Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.09% (W/V) sodium azide.

Storage Conditions

-20°C

ZADH2 Polyclonal Antibody - Protein Information**Name** PTGR3 ([HGNC:28697](#))**Synonyms** ZADH2**Function**

Functions as 15-oxo-prostaglandin 13-reductase and acts on 15-keto-PGE1, 15-keto-PGE2, 15-keto-PGE1-alpha and 15-keto-PGE2-alpha with highest efficiency towards 15-keto-PGE2-alpha. Overexpression represses transcriptional activity of PPARG and inhibits adipocyte differentiation.

Cellular Location



Peroxisome {ECO:0000250|UniProtKB:Q8BGC4}.

ZADH2 Polyclonal Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

ZADH2 Polyclonal Antibody - Images



ZADH2 Polyclonal Antibody - Background

Functions as 15-oxo-prostaglandin 13-reductase and acts on 15-keto-PGE1, 15-keto-PGE2, 15-keto-PGE1-alpha and 15-keto-PGE2-alpha with highest efficiency towards 15-keto-PGE2-alpha. Overexpression represses transcriptional activity of PPARG and inhibits adipocyte differentiation.