

ZNF397 Polyclonal Antibody
Catalog # AP73147**Specification**

ZNF397 Polyclonal Antibody - Product Information

Application	WB, IHC-P, IF
Primary Accession	Q8NF99
Reactivity	Human
Host	Rabbit
Clonality	Polyclonal

ZNF397 Polyclonal Antibody - Additional Information**Gene ID** 84307**Other Names**

ZNF397; ZNF47; ZSCAN15; Zinc finger protein 397; Zinc finger and SCAN domain-containing protein 15; Zinc finger protein 47

Dilution

WB~~Western Blot: 1/500 - 1/2000. Immunohistochemistry: 1/100 - 1/300. Immunofluorescence: 1/200 - 1/1000. ELISA: 1/5000. Not yet tested in other applications.

IHC-P~~N/A

IF~~1:50~200

Format

Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.09% (W/V) sodium azide.

Storage Conditions

-20°C

ZNF397 Polyclonal Antibody - Protein Information**Name** ZNF397**Synonyms** ZNF47, ZSCAN15**Function**

Isoform 3 acts as a DNA-dependent transcriptional repressor.

Cellular Location

[Isoform 1]: Nucleus.

Tissue Location


Expressed strongly in testis, moderately in skeletal muscle, pancreas and prostate, and weakly in heart, placenta, liver, kidney, spleen, thymus and small intestine

ZNF397 Polyclonal Antibody - Protocols


Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

ZNF397 Polyclonal Antibody - Images



Western Blot analysis of various cells using ZNF397 Polyclonal Antibody. Secondary antibody was diluted at 1:20000



Western Blot analysis of various cells using ZNF397 Polyclonal Antibody. Secondary antibody was diluted at 1:20000

ZNF397 Polyclonal Antibody - Background

Isoform 3 acts as a DNA-dependent transcriptional repressor.