

CHD1L Polyclonal Antibody
Catalog # AP73210**Specification****CHD1L Polyclonal Antibody - Product Information**

Application	WB
Primary Accession	Q86WJ1
Reactivity	Human, Mouse
Host	Rabbit
Clonality	Polyclonal

CHD1L Polyclonal Antibody - Additional Information**Gene ID** 9557**Other Names**

CHD1L; ALC1; Chromodomain-helicase-DNA-binding protein 1-like; Amplified in liver cancer protein 1

Dilution

WB~~Western Blot: 1/500 - 1/2000. ELISA: 1/20000. Not yet tested in other applications.

Format

Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.09% (W/V) sodium azide.

Storage Conditions

-20°C

CHD1L Polyclonal Antibody - Protein Information**Name** CHD1L {ECO:0000303|PubMed:34210977, ECO:0000312|HGNC:HGNC:1916}**Function**

ATP-dependent chromatin remodeler that mediates chromatin- remodeling following DNA damage (PubMed:19661379, PubMed:29220652, PubMed:29220653, PubMed:33357431, PubMed:34210977, PubMed:34486521, PubMed:34874266). Recruited to DNA damage sites through interaction with poly-ADP-ribose: specifically recognizes and binds histones that are poly-ADP-ribosylated on serine residues in response to DNA damage (PubMed:19661379, PubMed:29220652, PubMed:29220653, PubMed:34486521, PubMed:34874266).

Poly-ADP-ribose-binding activates the ATP-dependent chromatin remodeler activity, thereby regulating chromatin during DNA repair (PubMed:19661379, PubMed:29220652, PubMed:29220653, PubMed:34486521, PubMed:34874266). Catalyzes nucleosome sliding away from DNA breaks in an ATP-dependent manner (PubMed:19661379, PubMed:29220652, PubMed:29220653). Chromatin remodeling activity promotes PARP2 removal from chromatin (PubMed:33275888).

Cellular Location

Nucleus. Chromosome Note=Localizes at sites of DNA damage; recruited by histones H2B and H3 poly-ADP-ribosylated on 'Ser-6' and 'Ser-10', respectively (H2BS6ADPr and H3S10ADPr) by PARP1 or PARP2.

Tissue Location


Frequently overexpressed in hepatomacellular carcinomas.


CHD1L Polyclonal Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

CHD1L Polyclonal Antibody - Images





CHD1L Polyclonal Antibody - Background

DNA helicase which plays a role in chromatin-remodeling following DNA damage. Targeted to sites of DNA damage through interaction with poly(ADP-ribose) and functions to regulate chromatin during DNA repair. Able to catalyze nucleosome sliding in an ATP-dependent manner. Helicase activity is strongly stimulated upon poly(ADP-ribose)-binding.