

**GI Syn Polyclonal Antibody**  
**Catalog # AP73218****Specification****GI Syn Polyclonal Antibody - Product Information**

Application	WB
Primary Accession	<a href="#">P15104</a>
Reactivity	Human, Mouse, Rat
Host	Rabbit
Clonality	Polyclonal

**GI Syn Polyclonal Antibody - Additional Information****Gene ID** 2752**Other Names**

GLUL; GLNS; Glutamine synthetase; GS; Glutamate decarboxylase; Glutamate--ammonia ligase

**Dilution**

WB~~Western Blot: 1/500 - 1/2000. ELISA: 1/20000. Not yet tested in other applications.

**Format**

Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.09% (W/V) sodium azide.

**Storage Conditions**

-20°C

**GI Syn Polyclonal Antibody - Protein Information****Name** GLUL {ECO:0000303|PubMed:30158707, ECO:0000312|HGNC:HGNC:4341}**Function**

Glutamine synthetase that catalyzes the ATP-dependent conversion of glutamate and ammonia to glutamine (PubMed:<a href="http://www.uniprot.org/citations/16267323" target="\_blank">16267323</a>, PubMed:<a href="http://www.uniprot.org/citations/30158707" target="\_blank">30158707</a>, PubMed:<a href="http://www.uniprot.org/citations/36289327" target="\_blank">36289327</a>). Its role depends on tissue localization: in the brain, it regulates the levels of toxic ammonia and converts neurotoxic glutamate to harmless glutamine, whereas in the liver, it is one of the enzymes responsible for the removal of ammonia (By similarity). Plays a key role in ammonium detoxification during erythropoiesis: the glutamine synthetase activity is required to remove ammonium generated by porphobilinogen deaminase (HMBS) during heme biosynthesis to prevent ammonium accumulation and oxidative stress (By similarity). Essential for proliferation of fetal skin fibroblasts (PubMed:<a href="http://www.uniprot.org/citations/18662667" target="\_blank">18662667</a>). Independently of its glutamine synthetase activity, required for endothelial cell migration during vascular development: acts by regulating membrane localization and activation of the GTPase RHOJ, possibly by promoting RHOJ palmitoylation (PubMed:<a href="http://www.uniprot.org/citations/30158707" target="\_blank">30158707</a>). May act as a palmitoyltransferase for RHOJ: able to autopalmitylate and then transfer the palmitoyl group to

RHOJ (PubMed:<a href="http://www.uniprot.org/citations/30158707" target="\_blank">30158707</a>). Plays a role in ribosomal 40S subunit biogenesis (PubMed:<a href="http://www.uniprot.org/citations/26711351" target="\_blank">26711351</a>). Through the interaction with BEST2, inhibits BEST2 channel activity by affecting the gating at the aperture in the absence of intracellular L-glutamate, but sensitizes BEST2 to intracellular L-glutamate, which promotes the opening of BEST2 and thus relieves its inhibitory effect on BEST2 (PubMed:<a href="http://www.uniprot.org/citations/36289327" target="\_blank">36289327</a>).

#### Cellular Location

Cytoplasm, cytosol. Microsome {ECO:0000250|UniProtKB:P09606} Mitochondrion {ECO:0000250|UniProtKB:P09606}. Cell membrane; Lipid-anchor. Note=Mainly localizes in the cytosol, with a fraction associated with the cell membrane

#### Tissue Location


Expressed in endothelial cells.

### GI Syn Polyclonal Antibody - Protocols

Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

### GI Syn Polyclonal Antibody - Images



### GI Syn Polyclonal Antibody - Background

Glutamine synthetase that catalyzes the ATP-dependent conversion of glutamate and ammonia to glutamine (PubMed:30158707, PubMed:16267323). Its role depends on tissue localization: in the

brain, it regulates the levels of toxic ammonia and converts neurotoxic glutamate to harmless glutamine, whereas in the liver, it is one of the enzymes responsible for the removal of ammonia (By similarity). Essential for proliferation of fetal skin fibroblasts (PubMed:18662667). Independently of its glutamine synthetase activity, required for endothelial cell migration during vascular development: acts by regulating membrane localization and activation of the GTPase RHOJ, possibly by promoting RHOJ palmitoylation (PubMed:30158707). May act as a palmitoyltransferase for RHOJ: able to autopalmitylate and then transfer the palmitoyl group to RHOJ (PubMed:30158707).