

**Sox-17 Polyclonal Antibody**  
**Catalog # AP73238****Specification****Sox-17 Polyclonal Antibody - Product Information**

Application	WB
Primary Accession	<a href="#">Q9H6I2</a>
Reactivity	Human, Mouse
Host	Rabbit
Clonality	Polyclonal

**Sox-17 Polyclonal Antibody - Additional Information****Gene ID** 64321**Other Names**

SOX17; Transcription factor SOX-17

**Dilution**

WB~~Western Blot: 1/500 - 1/2000. ELISA: 1/20000. Not yet tested in other applications.

**Format**

Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.09% (W/V) sodium azide.

**Storage Conditions**

-20°C

**Sox-17 Polyclonal Antibody - Protein Information****Name** SOX17**Function**

Acts as a transcription regulator that binds target promoter DNA and bends the DNA. Binds to the sequences 5'-ACAAT-3' or 5'- ACAAAG-3'. Modulates transcriptional regulation via WNT3A. Inhibits Wnt signaling. Promotes degradation of activated CTNNB1. Plays a key role in the regulation of embryonic development. Required for normal development of the definitive gut endoderm. Required for normal looping of the embryonic heart tube. Plays an important role in embryonic and postnatal vascular development, including development of arteries. Plays an important role in postnatal angiogenesis, where it is functionally redundant with SOX18. Required for the generation and maintenance of fetal hematopoietic stem cells, and for fetal hematopoiesis. Probable transcriptional activator in the premeiotic germ cells.

**Cellular Location**

Nucleus {ECO:0000255|PROSITE-ProRule:PRU00267}.

**Tissue Location**

Expressed in adult heart, lung, spleen, testis, ovary, placenta, fetal lung, and kidney. In normal gastrointestinal tract, it is preferentially expressed in esophagus, stomach and small intestine than


in colon and rectum.

### **Sox-17 Polyclonal Antibody - Protocols**


Provided below are standard protocols that you may find useful for product applications.

- [Western Blot](#)
- [Blocking Peptides](#)
- [Dot Blot](#)
- [Immunohistochemistry](#)
- [Immunofluorescence](#)
- [Immunoprecipitation](#)
- [Flow Cytometry](#)
- [Cell Culture](#)

### **Sox-17 Polyclonal Antibody - Images**



Western Blot analysis of extracts from Jurkat cells, using Sox-17 Polyclonal Antibody.. Secondary antibody was diluted at 1:20000 cells nucleus extracted by Minute TM Cytoplasmic and Nuclear Fractionation kit (SC-003,Inventbiotech,MN,USA).



Western Blot analysis of extracts from Jurkat cells, using Sox-17 Polyclonal Antibody.. Secondary antibody was diluted at 1:20000 cells nucleus extracted by Minute TM Cytoplasmic and Nuclear Fractionation kit (SC-003,Inventbiotech,MN,USA).

### **Sox-17 Polyclonal Antibody - Background**

Acts as transcription regulator that binds target promoter DNA and bends the DNA. Binds to the sequences 5'- AACAAAT-3' or 5'-ACAAAG-3'. Modulates transcriptional regulation via WNT3A. Inhibits Wnt signaling. Promotes degradation of activated CTNNB1. Plays a key role in the regulation of embryonic development. Required for normal looping of the embryonic heart tube. Required for normal development of the definitive gut endoderm. Probable transcriptional activator in the premeiotic germ cells (By similarity).